arXiv:1601.03245v1 [math.GM] 12 Dec 2015

Equivalent of the elliptic function solutions of nonlinear differential equations

Dong-Hua Luo[1] and Cheng-Qun Pang[2,3*]
[1]*Shandong Communication Media College , Shandong Jinan 25020, China*
[2]*Department of Physics,Liupanshui Normal University Liupanshui, 553004, China*
[3]*School of Physical Science and Technology, Lanzhou University, Lanzhou 730000, China*

In this paper, we point out that many Jacobi elliptic function solutions to non-linear differential equation(NDE) can be transformed each other via the modulus and phase transformation of Jacobi elliptic function. Therefore these solutions are equivalent. We investigate equivalent of the Jacobi elliptic function solutions of Korteweg-de Vries (mKdV) equation as a example.

PACS numbers: 00A99; 97U60; 97U70

I. INTRODUCTION

In recent decades, nonlinear differential equation(NDE) become research focus, so the methods for solving them have been developed rapidly. Among them, a series Jacobi elliptic function solutions of nonlinear equations were given by the means of the Jacobi elliptic function expansion method[1] and its modified form[2,3] . In fact, many forms of Jacobi elliptic function solutions can be obtained from one of them using the transformation of the Jacobi elliptic function, therefore, they are equivalent. Further more, Jacobi elliptic function expansion method and its related correction method to reduce the number of expanded form, or expansion to certain restrictions, such as in Ref.[4] they have mentioned a few transformations of the Jacobi elliptic function, and used them to simplify the Jacobi elliptic function expansion. In Ref.[5], they pointed out that Jacobi elliptic transformation can be use to obtain other Jacobi elliptic functional solutions of NDE. In this paper, we point out that we should test wether the solutions are equivalent and we give a more systematic and comprehensive shown for transformation of the Jacobi elliptic function which can help us to test. Basing on any one of the known NDE's Jacobi elliptic function solutions, you can get other Jacobi elliptic function solutions by the transformation, of cause they are equivalent solutions. Finally, we investigate Jacobi elliptic function solutions to mKdV equation as a example.

The paper is organized as follows: In section II, we review the modulus and phase transformation of Jacobi elliptic function and give some new forms of them. A treatment for the Jacobi elliptic function solutions to mKdV equation as an example is shown in section III. In the last section, we give a brief conclusion.

II. JACOBI ELLIPTIC TRANSFORMATION

The definition and some properties of the Jacobi elliptic function can see appendix. Assuming $\mathrm{sn}(x+\phi|\lambda)$ can become $\mathrm{S}(x)$ by the module and phase transformation, at the same time $\mathrm{cn}(x+\phi|\lambda)$ and $\mathrm{dn}(x+\phi|\lambda)$ become $\mathrm{S}(x)$ and $\mathrm{D}(x)$ respectively, then S, C, and D meet the follow equations[8]

$$\frac{dS}{dx} = CD,\ S^2 + C^2 = 1,\ D^2 + \lambda S^2 = 1, \tag{1}$$

where ϕ can determined by equations

$$\mathrm{sn}(\phi|\lambda) = S(0),\ \mathrm{cn}(\phi|\lambda) = C(0),\ \mathrm{dn}(\phi|\lambda) = D(0). \tag{2}$$

If two Jacobi elliptic function solutions of a NDE satisfy (1) and (2), we can say that the two solutions are equivalence. By the means of appendix and (1) , we can get the follow expressions (some of them were given by Ref. [8])

$$\mathrm{sn}(\sqrt{k}x|\frac{1}{k}) = \sqrt{k}\mathrm{sn}(x|k), \tag{3}$$

$$\mathrm{cn}(\sqrt{k}x|\frac{1}{k}) = \mathrm{dn}(x|k), \tag{4}$$

*Electronic address: pangchq13@lzu.edu.cn

$$\mathrm{dn}(\sqrt{k}x|\frac{1}{k}) = \mathrm{cn}(x|k). \tag{5}$$

$$\mathrm{sn}\left(\sqrt{k}x + i\sqrt{k}K'(k) + \varphi|\frac{1}{k}\right) = (-1)^p \mathrm{ns}(x|k), \tag{6}$$

$$\mathrm{cn}\left(\sqrt{k}x + i\sqrt{k}K'(k) + \varphi|\frac{1}{k}\right) = (-1)^{p+q+1} i\mathrm{cs}(x|k), \tag{7}$$

$$\mathrm{dn}\left(\sqrt{k}x + i\sqrt{k}K'(k) + \varphi|\frac{1}{k}\right) = \frac{(-1)^{q+1}i}{\sqrt{k}}\mathrm{ds}(x|k), \tag{8}$$

$$\mathrm{sn}\left(\sqrt{k}x + \sqrt{k}K(k) + \varphi|\frac{1}{k}\right) = (-1)^p \sqrt{k} \cdot \mathrm{cd}(x|k), \tag{9}$$

$$\mathrm{cn}\left(\sqrt{k}x + \sqrt{k}K(k) + \varphi|\frac{1}{k}\right) = (-1)^{p+q} \sqrt{k'} \cdot \mathrm{nd}(x|k), \tag{10}$$

$$\mathrm{dn}\left(\sqrt{k}x + \sqrt{k}K(k) + \varphi|\frac{1}{k}\right) = (-1)^{q+1} \sqrt{k'}\mathrm{sd}(x|k), \tag{11}$$

$$\mathrm{sn}\left(\sqrt{k}x + i\sqrt{k}K'(k) + \sqrt{k}K(k) + \varphi|\frac{1}{k}\right) = (-1)^p \cdot \mathrm{dc}(x|k), \tag{12}$$

$$\mathrm{cn}\left(\sqrt{k}x + i\sqrt{k}K'(k) + \sqrt{k}K(k) + \varphi|\frac{1}{k}\right) = (-1)^{p+q} i\sqrt{k'} \cdot \mathrm{cs}(x|k), \tag{13}$$

$$\mathrm{dn}\left(\sqrt{k}x + i\sqrt{k}K'(k) + \sqrt{k}K(k) + \varphi|\frac{1}{k}\right) = (-1)^{q+1} i\sqrt{\frac{k'}{k}}\mathrm{nc}(x|k), \tag{14}$$

where $\varphi = 2pK(1/k) + 2iqK(1/k)'$.

$$\mathrm{sn}(ix|k') = i \cdot \mathrm{sc}(x|k), \tag{15}$$

$$\mathrm{cn}(ix|k') = \mathrm{nc}(x|k), \tag{16}$$

$$\mathrm{dn}(ix|k') = \mathrm{dc}(x|k), \tag{17}$$

$$\mathrm{sn}(i\sqrt{k}x| - \frac{k'}{k}) = i\sqrt{k} \cdot \mathrm{sd}(x|k), \tag{18}$$

$$\mathrm{cn}(i\sqrt{k}x| - \frac{k'}{k}) = \mathrm{nd}(x|k), \tag{19}$$

$$\mathrm{dn}(i\sqrt{k}x| - \frac{k'}{k}) = \mathrm{cd}(x|k), \tag{20}$$

$$\mathrm{sn}(\sqrt{k'}x| - \frac{k}{k'}) = \sqrt{k'}\mathrm{sd}(x|k), \tag{21}$$

$$\mathrm{cn}(\sqrt{k'}x| - \frac{k}{k'}) = \mathrm{cd}(x|k), \tag{22}$$

$$\mathrm{dn}(\sqrt{k'}x|-\frac{k}{k'}) = \mathrm{nd}(x|k), \tag{23}$$

$$\mathrm{sn}(i\sqrt{k'}x|\frac{1}{k'}) = i\sqrt{k'}\mathrm{sc}(x|k), \tag{24}$$

$$\mathrm{cn}(i\sqrt{k'}x|\frac{1}{k'}) = \mathrm{dc}(x|k), \tag{25}$$

$$\mathrm{dn}(i\sqrt{k'}x|\frac{1}{k'}) = \mathrm{nc}(x|k), \tag{26}$$

$$\mathrm{sn}\left(\left(\sqrt{k'}-\varepsilon\right)x|\left(\frac{\sqrt{k'}+\varepsilon}{\sqrt{k'}-\varepsilon}\right)^2\right) = \left(\sqrt{k'}-\varepsilon\right)\mathrm{cd}(x|k)\cdot\mathrm{sn}(x|k), \tag{27}$$

where $\varepsilon^2 = 1$,

$$\mathrm{cn}\left(\left(\sqrt{k'}-\varepsilon\right)x|\left(\frac{\sqrt{k'}+\varepsilon}{\sqrt{k'}-\varepsilon}\right)^2\right) = \left(\varepsilon\sqrt{k'}-1\right)\mathrm{sd}(x|k)\mathrm{sn}(x|k)+\mathrm{nd}(x|k), \tag{28}$$

$$\mathrm{dn}\left(\left(\sqrt{k'}-\varepsilon\right)x|\left(\frac{\sqrt{k'}+\varepsilon}{\sqrt{k'}-\varepsilon}\right)^2\right) = \frac{1-\left[\mathrm{dn}(x|k)^2+1\right]\mathrm{sn}(x|k)^2}{\mathrm{dn}(x|k)\left[\left(\varepsilon\sqrt{k'}-1\right)\mathrm{sn}(x|k))^2+1\right]}, \tag{29}$$

$$\mathrm{sn}\left(\frac{1}{2}\left(\varepsilon\sqrt{k'}+1\right)x+\phi|\left(\frac{1-\varepsilon\sqrt{k'}}{1+\varepsilon\sqrt{k'}}\right)^2\right) = \left(\varepsilon+\sqrt{k'}\right)\cdot\frac{\mathrm{cn}(x|k)}{\sqrt{k'}-\mathrm{dn}(x|k)}, \tag{30}$$

$$\mathrm{cn}\left(\frac{1}{2}\left(\varepsilon\sqrt{k'}+1\right)x+\phi|\left(\frac{1-\varepsilon\sqrt{k'}}{1+\varepsilon\sqrt{k'}}\right)^2\right) = i\varepsilon\sqrt{\frac{2}{k}}\sqrt[4]{k'}\frac{\sqrt{[\mathrm{dn}(x|k)+\varepsilon]\left\{\left(\sqrt{k'}+\varepsilon\right)[\mathrm{dn}(x|k)-\varepsilon]+k\right\}}}{\mathrm{dn}(x|k)-\sqrt{k'}}, \tag{31}$$

$$\mathrm{dn}\left(\frac{1}{2}\left(\varepsilon\sqrt{k'}+1\right)x+\phi|\left(\frac{1-\varepsilon\sqrt{k'}}{1+\varepsilon\sqrt{k'}}\right)^2\right) = \frac{i\sqrt{2k}\sqrt[4]{k'}\mathrm{sn}(x|k)}{\sqrt{[\mathrm{dn}(x|k)+\varepsilon]\left\{\left(\sqrt{k'}+\varepsilon\right)[\mathrm{dn}(x|k)-\varepsilon]+k\right\}}}, \tag{32}$$

where $\phi = -\varepsilon K + \frac{1}{2}(1+\varepsilon)iK'$,

$$\mathrm{sn}\left(\frac{i}{2}\left(\sqrt{k}+\varepsilon\right)x+\phi|\left(\frac{\sqrt{k}-\varepsilon}{\sqrt{k}+\varepsilon}\right)^2\right) = \varepsilon\sqrt{\frac{\varepsilon+\sqrt{k}}{\varepsilon-\sqrt{k}}}\frac{\mathrm{dn}(x|k)}{\sqrt{k}\mathrm{sn}(x|k)-1}, \tag{33}$$

$$\mathrm{cn}\left(\frac{i}{2}\left(\sqrt{k}+\varepsilon\right)x+\phi|\left(\frac{\sqrt{k}-\varepsilon}{\sqrt{k}+\varepsilon}\right)^2\right) = \frac{\varepsilon\sqrt{\left(\varepsilon+\sqrt{k}\right)\mathrm{dn}(x|k)^2+\left(\sqrt{k}-\varepsilon\right)\left[\sqrt{k}\mathrm{sn}(x|k)-1\right]^2}}{\sqrt{\sqrt{k}-\varepsilon}\left(1-\sqrt{k}\mathrm{sn}(x|k)\right)}, \tag{34}$$

$$\mathrm{dn}\left(\frac{i}{2}\left(\sqrt{k}+\varepsilon\right)x+\phi|\left(\frac{\sqrt{k}-\varepsilon}{\sqrt{k}+\varepsilon}\right)^2\right) = \frac{\mathrm{cn}(x|k)}{\varepsilon\sqrt{k}+1}\sqrt{\frac{k\left(\sqrt{k}+\varepsilon\right)}{\left(\varepsilon+\sqrt{k}\right)\mathrm{dn}(x|k)^2+\left(\sqrt{k}-\varepsilon\right)\left[\sqrt{k}\mathrm{sn}(x|k)-1\right]^2}}, \tag{35}$$

where $\phi = \varepsilon\left(-K + \frac{iK'}{2}\right)$,

$$\mathrm{sn}\left(\frac{1}{2}\left(i\varepsilon\sqrt{k'} + \sqrt{k}\right)x|\left(-\sqrt{k}\varepsilon + i\sqrt{1-k}\right)^4\right) = \left(i\varepsilon\sqrt{k'} + \sqrt{k}\right)\frac{\mathrm{sn}(x|k)}{1+\mathrm{cn}(x|k)}, \tag{36}$$

$$\mathrm{cn}\left(\frac{1}{2}\left(i\varepsilon\sqrt{k'} + \sqrt{k}\right)x|\left(-\sqrt{k}\varepsilon + i\sqrt{1-k}\right)^4\right) = \sqrt{\frac{2\left(\sqrt{k'} - i\varepsilon\sqrt{k}\right)\left[\sqrt{k'} + i\varepsilon\sqrt{k}\mathrm{cn}(x|k)\right]}{1+\mathrm{cn}(x|k)}}, \tag{37}$$

$$\mathrm{dn}\left(\frac{1}{2}\left(i\varepsilon\sqrt{k'} + \sqrt{k}\right)x|\left(-\sqrt{k}\varepsilon + i\sqrt{1-k}\right)^4\right) = \frac{\sqrt{2}\mathrm{dn}(x|k)}{\sqrt{\left(\sqrt{k'} - i\varepsilon\sqrt{k}\right)\left[\sqrt{k'} + i\varepsilon\sqrt{k}\mathrm{cn}(x|k)\right][1+\mathrm{cn}(x|k)]}}, \tag{38}$$

$$\mathrm{sn}\left(2\sqrt{\varepsilon\sqrt{k}}x + \phi|\frac{\varepsilon\left(\varepsilon + \sqrt{k}\right)^2}{4\sqrt{k}}\right) = \frac{i}{\left(\varepsilon + \sqrt{k}\right)}\left(\varepsilon\sqrt{k}\mathrm{sn}(x|k) + \mathrm{ns}(x|k)\right), \tag{39}$$

$$\mathrm{cn}\left(2\sqrt{\varepsilon\sqrt{k}}x + \phi|\frac{\varepsilon\left(\varepsilon + \sqrt{k}\right)^2}{4\sqrt{k}}\right) = \frac{-i}{\left(1 + \varepsilon\sqrt{k}\right)}\left(\varepsilon\sqrt{k}\mathrm{cs}(x|k)\mathrm{dn}(x|k)\right), \tag{40}$$

$$\mathrm{dn}\left(2\sqrt{\varepsilon\sqrt{k}}x + \phi|\frac{\varepsilon\left(\varepsilon + \sqrt{k}\right)^2}{4\sqrt{k}}\right) = \frac{\sqrt{-\varepsilon}}{2\sqrt[4]{k}}\left(\sqrt{k}\mathrm{sn}(x|k) - \varepsilon\mathrm{ns}(x|k)\right), \tag{41}$$

where $\phi = \varepsilon i K'$,

$$\mathrm{sn}\left(2\sqrt{\varepsilon}\sqrt[4]{-kk'}x + \phi|\frac{\varepsilon\left(\sqrt{-k'} - \varepsilon\sqrt{k}\right)^2}{4\sqrt{-kk'}}\right) = \left(\varepsilon\sqrt{-k'} + \sqrt{k}\right)\left(\sqrt{k}\mathrm{cn}(x|k) - \varepsilon\sqrt{-k'}\mathrm{nc}(x|k)\right), \tag{42}$$

$$\mathrm{cn}\left(2\sqrt{\varepsilon}\sqrt[4]{-kk'}x + \phi|\frac{\varepsilon\left(\sqrt{-k'} - \varepsilon\sqrt{k}\right)^2}{4\sqrt{-kk'}}\right) = -\left(\sqrt{k} + \varepsilon\sqrt{-k'}\right)\mathrm{sn}(x|k)\mathrm{dc}(x|k), \tag{43}$$

$$\mathrm{dn}\left(2\sqrt{\varepsilon}\sqrt[4]{-kk'}x + \phi|\frac{\varepsilon\left(\sqrt{-k'} - \varepsilon\sqrt{k}\right)^2}{4\sqrt{-kk'}}\right) = \frac{\varepsilon^{1/2}\sqrt{k}}{2\sqrt[4]{-k'k}}\left(\mathrm{cn}(x|k) + \varepsilon\sqrt{\frac{-k'}{k}}\mathrm{nc}(x|k)\right), \tag{44}$$

where $\phi = \varepsilon K$,

$$\mathrm{sn}\left(2\sqrt{\varepsilon\sqrt{k'}}x + \phi|\frac{-\varepsilon\left(\sqrt{k'} - \varepsilon\right)^2}{4\sqrt{k'}}\right) = \frac{-1}{\varepsilon - \sqrt{k'}}\left[\varepsilon\mathrm{dn}(x|k) - \sqrt{k'}\mathrm{nd}(x|k)\right], \tag{45}$$

$$\mathrm{cn}\left(2\sqrt{\varepsilon\sqrt{k'}}xx + \phi|\frac{-\varepsilon\left(\sqrt{k'} - \varepsilon\right)^2}{4\sqrt{k'}}\right) = \left(\sqrt{k'} + \varepsilon\right)\mathrm{sn}(x|k)\mathrm{cd}(x|k), \tag{46}$$

$$\mathrm{dn}\left(2\sqrt{\varepsilon\sqrt{k'}}x + \phi|\frac{-\varepsilon\left(\sqrt{k'} - \varepsilon\right)^2}{4\sqrt{k'}}\right) = \frac{\varepsilon^{3/2}}{2\sqrt[4]{k'}}\left[\varepsilon\mathrm{dn}(x|k) + \sqrt{k'}\mathrm{nd}(x|k)\right], \tag{47}$$

where $\phi = -K$,

$$\mathrm{sn}\left(\frac{1}{2}i\varepsilon\left(\sqrt[4]{k}+\varepsilon^{3/2}\right)^2 x+\phi|\left(\frac{\sqrt[4]{k}-\varepsilon^{3/2}}{\sqrt[4]{k}+\varepsilon^{3/2}}\right)^4\right)=\frac{-\left(\sqrt[4]{k}+\varepsilon^{3/2}\right)}{\sqrt[4]{k}+\varepsilon^{3/2}}\frac{\varepsilon^{3/2}+\sqrt[4]{k}\mathrm{sn}(x|k)}{\varepsilon^{3/2}-\sqrt[4]{k}\mathrm{sn}(x|k)}, \tag{48}$$

$$\mathrm{cn}\left(\frac{1}{2}i\varepsilon\left(\sqrt[4]{k}+\varepsilon^{3/2}\right)^2 x+\phi|\left(\frac{\sqrt[4]{k}-\varepsilon^{3/2}}{\sqrt[4]{k}+\varepsilon^{3/2}}\right)^4\right)=\frac{2i\varepsilon^{3/4}\sqrt[8]{k}\sqrt{(1+\mathrm{sn}(x|k))\left(\varepsilon+\sqrt{k}\mathrm{sn}(x|k)\right)}}{\left(\sqrt[4]{k}-\varepsilon^{3/2}\right)\left(\varepsilon^{3/2}-\sqrt[4]{k}\mathrm{sn}(x|k)\right)}, \tag{49}$$

$$\mathrm{dn}\left(\frac{1}{2}i\varepsilon\left(\sqrt[4]{k}+\varepsilon^{3/2}\right)^2 x+\phi|\left(\frac{\sqrt[4]{k}-\varepsilon^{3/2}}{\sqrt[4]{k}+\varepsilon^{3/2}}\right)^4\right)=\frac{2\varepsilon^{-1/4}\sqrt[8]{k}\mathrm{cn}(x|k)\mathrm{dn}(x|k)}{\left(\sqrt[4]{k}+\varepsilon^{3/2}\right)\left(\varepsilon^{3/2}-\sqrt[4]{k}\mathrm{sn}(x|k)\right)\sqrt{(1+\mathrm{sn}(x|k))\left(\varepsilon+\sqrt{k}\mathrm{sn}(x|k)\right)}}, \tag{50}$$

where $\phi = \frac{\varepsilon+1}{2}K + \frac{iK'}{2}$,

$$\mathrm{sn}\left(\frac{1}{2}i\left((-\varepsilon)^{3/2}\sqrt[4]{k'}+\sqrt[4]{-k}\right)^2 x+\phi|\left(\frac{\sqrt[4]{-k}-(-\varepsilon)^{3/2}\sqrt[4]{k'}}{\sqrt[4]{-k}+(-\varepsilon)^{3/2}\sqrt[4]{k'}}\right)^4\right)=-\frac{\left(\sqrt[4]{-k}+(-\varepsilon)^{3/2}\sqrt[4]{k'}\right)}{\sqrt[4]{-k}-(-\varepsilon)^{3/2}\sqrt[4]{k'}}\frac{\left((-\varepsilon)^{3/2}\sqrt[4]{k'}+\sqrt[4]{-k}\mathrm{cn}(x|k)\right)}{\left((-\varepsilon)^{3/2}\sqrt[4]{k'}-\sqrt[4]{-k}\mathrm{cn}(x|k)\right)}, \tag{51}$$

$$\begin{aligned}&\mathrm{cn}\left(\frac{1}{2}i\left((-\varepsilon)^{3/2}\sqrt[4]{k'}+\sqrt[4]{-k}\right)^2 x+\phi|\left(\frac{\sqrt[4]{-k}-(-\varepsilon)^{3/2}\sqrt[4]{k'}}{\sqrt[4]{-k}+(-\varepsilon)^{3/2}\sqrt[4]{k'}}\right)^4\right)\\&=\frac{2i\sqrt[8]{-kk'}\sqrt{(2k-\varepsilon)^2}\sqrt{(\mathrm{cn}(x|k)+1)\left(\sqrt{k}\mathrm{cn}(x|k)-\varepsilon\sqrt{k'}\right)}}{(-\varepsilon)^{1/4}(2k-\varepsilon)\left(\sqrt[4]{-k}-(-\varepsilon)^{3/2}\sqrt[4]{k'}\right)\left(\sqrt[4]{-k}\mathrm{cn}(x|k)-(-\varepsilon)^{3/2}\sqrt[4]{k'}\right)},\end{aligned} \tag{52}$$

$$\begin{aligned}&\mathrm{dn}\left(\frac{1}{2}i\left((-\varepsilon)^{3/2}\sqrt[4]{k'}+\sqrt[4]{-k}\right)^2 x+\phi|\left(\frac{\sqrt[4]{-k}-(-\varepsilon)^{3/2}\sqrt[4]{k'}}{\sqrt[4]{-k}+(-\varepsilon)^{3/2}\sqrt[4]{k'}}\right)^4\right)\\&=\frac{1}{\sqrt{(\mathrm{cn}(x|k)+1)\left(\sqrt{-k}\mathrm{cn}(x|k)-\varepsilon\sqrt{k'}\right)}}\frac{-2(-\varepsilon)^{-1/4}\sqrt[8]{-kk'}\sqrt{(2k-1)^2}\mathrm{sn}(x|k)\mathrm{dn}(x|k)}{(2k-\varepsilon)\left(\sqrt[4]{-k}+(-\varepsilon)^{3/2}\sqrt[4]{k'}\right)\left(\sqrt[4]{-k}\mathrm{cn}(x|k)-(-\varepsilon)^{3/2}\sqrt[4]{k'}\right)},\end{aligned} \tag{53}$$

where $\phi = -K - iK'$,

$$\mathrm{sn}\left(\frac{1}{2}\left(1+\varepsilon^{3/2}\sqrt[4]{k'}\right)^2 x+\phi|\left(\frac{1+\varepsilon^{3/2}\sqrt[4]{k'}}{1-\varepsilon^{3/2}\sqrt[4]{k'}}\right)^4\right)=-\varepsilon\frac{\left(1+\varepsilon^{3/2}\sqrt[4]{k'}\right)}{\left(1-\varepsilon^{3/2}\sqrt[4]{k'}\right)}\frac{\mathrm{dn}(x|k)+\varepsilon^{3/2}\sqrt[4]{k'}}{\mathrm{dn}(x|k)-\varepsilon^{3/2}\sqrt[4]{k'}}, \tag{54}$$

$$\mathrm{cn}\left(\frac{1}{2}\left(1+\varepsilon^{3/2}\sqrt[4]{k'}\right)^2 x+\phi|\left(\frac{1+\varepsilon^{3/2}\sqrt[4]{k'}}{1-\varepsilon^{3/2}\sqrt[4]{k'}}\right)^4\right)=\frac{2i\varepsilon^{3/4}\sqrt[8]{k'}\sqrt{(1+\mathrm{dn}(x|k))\left(\varepsilon\sqrt{k'}+\mathrm{dn}(x|k)\right)}}{\left(1-\varepsilon^{3/2}\sqrt[4]{k'}\right)\left(\mathrm{dn}(x|k)-\varepsilon^{3/2}\sqrt[4]{k'}\right)}, \tag{55}$$

$$\mathrm{dn}\left(\tfrac{1}{2}\left(1+\varepsilon^{3/2}\sqrt[4]{k'}\right)^2 x+\phi|\left(\tfrac{1+\varepsilon^{3/2}\sqrt[4]{k'}}{1-\varepsilon^{3/2}\sqrt[4]{k'}}\right)^4\right)=\frac{2ik\varepsilon^{-1/4}k'^{1/8}\mathrm{sn}(x|k)\mathrm{cn}(x|k)}{\left(1+\varepsilon^{3/2}\sqrt[4]{k'}\right)\left(\mathrm{dn}(x|k)-\varepsilon^{3/2}\sqrt[4]{k'}\right)\sqrt{(1+\mathrm{dn}(x|k))\left(\varepsilon\sqrt{k'}+\mathrm{dn}(x|k)\right)}}, \tag{56}$$

where $\phi = -K - iK'$.

III. INVESTIGATION OF JACOBI ELLIPTIC FUNCTION SOLUTIONS TO MKDV

In this section, we will apply our transform to NDE. We take the mKdV equation for example. The mKdV equation reads

$$u_t + 6\nu u^2 u_x + u_{xxx} = 0, \tag{57}$$

where $\nu^2 = 1$, $\nu = 1$ for positive mKdV(p-mKdV) equation, and $\nu = -1$ for negative mKdV(n-mKdV) equation. If we use the relation:

$$u = 2\sqrt{\nu}\frac{\partial \tan^{-1}(\phi)}{\partial x}, \tag{58}$$

Thus substitution of (58) in (57) gives:

$$\left(1+\phi^2\right)(\phi_t + \phi_{xxx}) + 6\phi_x\left(\phi^2 - \phi\phi_{xx}\right) = 0, \tag{59}$$

where ϕ is the function of x and t. Ref. [5,9–11] solved the equation (59), and got some Jacobi elliptic function solutions. Fu[11] obtained p-mKdV's more binary Jacobi elliptic function solutions using a systematical way, and pointed out the conditions for the p-mKdV's binary Jacobi elliptic function solutions existing. We will discuss the conditions for real solution of both p-mKdV and n-mKdV. As all we known:

$$\tanh^{-1}(i\phi) = i\tan^{-1}(\phi),\ i\tanh^{-1}(\phi) = \tan^{-1}(i\phi), \tag{60}$$

combining (58), we can find that if ϕ is imaginary number, then the solution of n-mKdV will be real. Therefor if considering n-mKdV and p-mKdV's real solutions at the same time, we can treat ϕ^2 as a real. So when we known a solution of (59) given by Ref. [5], we can get a lot of Jacobi elliptic function solutions of n-mKdV and p-mKdV. Such as

$$\phi_0 = \frac{ia\sqrt{k}\sigma}{c}\mathrm{sn}\left(\xi|k\right)\mathrm{sn}\left(\eta|m\right), \tag{61}$$

where

$$\begin{aligned} \frac{c^4}{a^4} &= \frac{k}{m}, \\ b &= a\left((k+1)a^2 + 3c^2(m+1)\right), \\ \frac{d}{c} &= 3(k+1)a^2 + c^2(m+1), \end{aligned} \tag{62}$$

and

$$\xi = ax + bt + a_0,\ \eta = cx + dt + c_0,\ \sigma^2 = 1. \tag{63}$$

By the means of (21), using the transform $\xi \to \sqrt{k'}\xi,\ \eta \to \sqrt{m'}\eta,\ k \to \frac{k}{-k'},\ m \to \frac{m}{-m'}$, we can get

$$\phi_1 = \sigma\frac{a}{c}\sqrt{k'k}\,\mathrm{sd}(\mathrm{a0} + bt + ax|k)\mathrm{sd}(\mathrm{c0} + dt + cx|m), \tag{64}$$

where

$$\frac{c^4}{a^4} = \frac{k'k}{m'm},\ \frac{b}{a}(1-2k)a^2 + 3c^2(1-2m),\ \frac{d}{c} = (3-6k)a^2 + c^2(1-2m). \tag{65}$$

The other Jacobi elliptic function solutions of mKdV equation are list in Table I. So these solutions of mKdV equation are equivalent.

TABLE I: Jacobi elliptic function solutions of mKdV

number $2\sqrt{\nu}\frac{\partial\tan^{-1}}{\partial x}$		$\frac{c^4}{a^4}$, $\frac{b}{a}$, $\frac{d}{c}$
1	$\frac{i\sigma a\ \sqrt{k}\mathrm{sc}(\eta\|m)\mathrm{sn}(\xi\|k)}{c}$	$\frac{k}{m'}, (k+1)a^2 + 3c^2(m-2), 3(k+1)a^2 + c^2(m-2)$
2	$\frac{\mathrm{i}\sigma a\ \sqrt{k'}\mathrm{sc}(\eta\|m)\mathrm{sc}(\xi\|k)}{c}$	$\frac{k'}{m'}, (k-2)a^2 + 3c^2(m-2), 3(k-2)a^2 + c^2(m-2)$
3	$\frac{a\sigma\ \sqrt{k}\ \sqrt{-m}\ \mathrm{sd}(\eta\|m)\mathrm{sn}(\xi\|k)}{c\sqrt{m}}$	$-\frac{k}{mm'}, (k+1)a^2 + 3c^2(1-2m), 3(k+1)a^2 + c^2(1-2m)$
4	$\frac{\sigma a\ \sqrt{k'}m^{5/2}\mathrm{sc}(\xi\|k)\mathrm{sd}(\eta\|m)}{c(-m)^{5/2}}$	$-\frac{k'}{mm'}, (k-2)a^2 + 3c^2(1-2m), 3(k-2)a^2 + c^2(1-2m)$
5	$\frac{\sigma a\ \sqrt{kk'}\ \mathrm{sd}(\eta\|m)\mathrm{sd}(\xi\|k)}{c}$	$\frac{kk'}{mm'}, (1-2k)a^2 + 3c^2(1-2m), (3-6k)a^2 + c^2(1-2m)$
6	$\frac{ia\ \sqrt{k}\sigma\mathrm{cn}(\eta\|m)\mathrm{sn}(\eta\|m)\mathrm{sn}(\xi\|k)}{c\mathrm{dn}(\eta\|m)}$	$\frac{k}{m^2}, a^2(k+1) - 6c^2(m-2), 3a^2(k+1) - 2c^2(m-2)$
7	$\frac{i\sigma a\ \sqrt{k'}\mathrm{cn}(\eta\|m)\mathrm{sc}(\xi\|k)\mathrm{sn}(\eta\|m)}{c\mathrm{dn}(\eta\|m)}$	$\frac{k'}{m^2}, a^2(k-2) - 6c^2(m-2), 3a^2(k-2) - 2c^2(m-2)$
8	$\frac{a\sigma\ \sqrt{kk'}\mathrm{cn}(\eta\|m)\mathrm{sd}(\xi\|k)\mathrm{sn}(\eta\|m)}{c\mathrm{dn}(\eta\|m)}$	$-\frac{kk'}{m^2}, a^2(1-2k) - 6c^2(m-2), a^2(3-6k) - 2c^2(m-2)$

continue I

number $2\sqrt{\nu}\frac{\partial\tan^{-1}}{\partial x}$		$\frac{c^4}{a^4}, \frac{b}{a}, \frac{d}{c}$
9	$\frac{iak\sigma\mathrm{cn}(\eta\|m)\mathrm{cn}(\xi\|k)\mathrm{sn}(\eta\|m)\mathrm{sn}(\xi\|k)}{\mathrm{cdn}(\eta\|m)\mathrm{dn}(\xi\|k)}$	$\frac{k^2}{m^2}, -2(k-2)a^2-6c^2(m-2), -6(k-2)a^2-2c^2(m-2)$
10	$\frac{2ia\sqrt{k}\sigma\mathrm{cn}(\eta\|m)\mathrm{sn}(\xi\|k)\sigma_m}{c\left(\sqrt{m'}-\mathrm{dn}(\eta\|m)\right)}$	$\frac{16k}{m^2}, a^2(k+1)-\frac{3}{2}c^2(m-2), 3a^2(k+1)-\frac{c^2}{2}(m-2)$
11	$\frac{2iak\sigma\sigma_m\sqrt{k'}\left(\sigma_m+\sqrt{m'}\right)\mathrm{cn}(\eta\|m)\mathrm{sc}(\xi\|k)}{c\left(\sqrt{m'}-\mathrm{dn}(\eta\|m)\right)}$	$\frac{16k'}{m^2}, \frac{1}{2}\left(2a^2(k-2)-3c^2(m-2)\right), 3a^2(k-2)-\frac{c^2}{2}(m-2)$
12	$\frac{2a\sigma\sigma_m\sqrt{kk'}\left(\sigma_m+\sqrt{m'}\right)\mathrm{cn}(\eta\|m)\mathrm{sd}(\xi\|k)}{c\left(\sqrt{m'}-\mathrm{dn}(\eta\|m)\right)}$	$-\frac{16kk'}{m^2}, a^2(1-2k)-\frac{3}{2}c^2(m-2), a^2(3-6k)-\frac{1}{2}c^2(m-2)$
13	$\frac{2iak\sigma\sigma_m\left(\sigma_m+\sqrt{m'}\right)\mathrm{cn}(\eta\|m)\mathrm{cn}(\xi\|k)\mathrm{sn}(\xi\|k)}{c\mathrm{dn}(\xi\|k)\left(\sqrt{m'}-\mathrm{dn}(\eta\|m)\right)}$	$\frac{16k^2}{m^2}, \frac{1}{2}\left(-4(k-2)a^2-3c^2(m-2)\right), -6(k-2)a^2-\frac{1}{2}c^2(m-2)$
14	$\frac{iak\sigma\sigma_k\sigma_m\left(\sigma_m+\sqrt{m'}\right)\mathrm{cn}(\eta\|m)\mathrm{cn}(\xi\|k)}{c\left(\sqrt{k'}-\mathrm{dn}(\xi\|k)\right)\left(\sqrt{m'}-\mathrm{dn}(\eta\|m)\right)}$	$\frac{k^2}{m^2}, \frac{1}{2}\left(a^2(2-k)-3c^2(m-2)\right), -\frac{3}{2}(k-2)a^2-\frac{1}{2}c^2(m-2)$
15	$\frac{2a\sqrt{k}\sigma\sigma_m\mathrm{dn}(\eta\|m)\mathrm{sn}(\xi\|k)}{c\sqrt{m'}\left(\sqrt{m}\mathrm{sn}(\eta\|m)-1\right)}$	$\frac{16k}{m'^2}, a^2(k+1)-\frac{3}{2}c^2(m+1), 3a^2(k+1)-\frac{c^2}{2}(m+1)$
16	$\frac{2a\sigma\sigma_m\sqrt{k'}\mathrm{dn}(\eta\|m)\mathrm{sc}(\xi\|k)}{c\sqrt{m'}\left(\sqrt{m}\mathrm{sn}(\eta\|m)-1\right)}$	$\frac{16k'}{m'^2}, a^2(k-2)-\frac{3}{2}c^2(m+1), 3a^2(k-2)-\frac{1}{2}c^2(m+1)$
17	$\frac{2ia\sigma\sigma_m\sqrt{kk'}\mathrm{dn}(\eta\|m)\mathrm{sd}(\xi\|k)}{c\sqrt{m'}\left(\sqrt{m}\mathrm{sn}(\eta\|m)-1\right)}$	$-\frac{16kk'}{m'^2}, a^2(1-2k)-\frac{3}{2}c^2(m+1), a^2(3-6k)-\frac{1}{2}c^2(m+1)$
18	$\frac{2ak\sigma\sigma_m\mathrm{cn}(\xi\|k)\mathrm{dn}(\eta\|m)\mathrm{sn}(\xi\|k)}{c\sqrt{m'}\mathrm{dn}(\xi\|k)\left(\sqrt{m}\mathrm{sn}(\eta\|m)-1\right)}$	$\frac{16k^2}{m'^2}, -2(k-2)a^2-\frac{3}{2}c^2(m+1), -6(k-2)a^2-\frac{1}{2}c^2(m+1)$
19	$\frac{ak\sigma\sigma_k\sigma_m\mathrm{cn}(\xi\|k)\mathrm{dn}(\eta\|m)}{c\sqrt{m'}\left(\sqrt{m}\mathrm{sn}(\eta\|m)-1\right)\left(\sqrt{k'}-\mathrm{dn}(\xi\|k)\right)}$	$\frac{k^2}{m'^2}, \frac{1}{2}a^2(2-k)-\frac{3}{2}c^2(m+1), -\frac{3}{2}(k-2)a^2-\frac{c^2}{2}(m+1)$
20	$\frac{ia\sigma\sigma_k\sigma_m\sqrt{k'}\mathrm{dn}(\eta\|m)\mathrm{dn}(\xi\|k)}{c\sqrt{m'}\left(\sqrt{m}\mathrm{sn}(\eta\|m)-1\right)\left(\sqrt{k}\mathrm{sn}(\xi\|k)-1\right)}$	$\frac{k'^2}{m'^2}, \frac{1}{2}a^2(-k-1)-\frac{3}{2}c^2(m+1), -\frac{3}{2}(k+1)a^2-\frac{c^2}{2}(m+1)$
21	$\frac{2a\sigma\sigma_m\left(\sqrt{m}\sigma_m+i\sqrt{m'}\right)\sqrt{k}\mathrm{sn}(\eta\|m)\mathrm{sn}(\xi\|k)}{c(\mathrm{cn}(\eta\|m)+1)\left(\sigma_m\sqrt{m'}-i\sqrt{m}\right)}$	$16k, (k+1)a^2+\frac{3}{2}c^2(2m-1), 3(k+1)a^2+\frac{1}{2}c^2(2m-1)$
22	$\frac{2a\sigma\sigma_m\sqrt{k'}\left(\sqrt{m}\sigma_m+i\sqrt{m'}\right)\mathrm{sc}(\xi\|k)\mathrm{sn}(\eta\|m)}{c\left(\sigma_m\sqrt{m'}-i\sqrt{m}\right)(\mathrm{cn}(\eta\|m)+1)}$	$16k', (k-2)a^2+\frac{3}{2}c^2(2m-1), 3(k-2)a^2+\frac{1}{2}c^2(2m-1)$
23	$\frac{2a\sigma\sigma_m\sqrt{kk'}\left(\sqrt{m}\sigma_m+i\sqrt{m'}\right)\mathrm{sd}(\xi\|k)\mathrm{sn}(\eta\|m)}{c\left(i\sqrt{m'}\sigma_m+\sqrt{m}\right)(\mathrm{cn}(\eta\|m)+1)}$	$-16kk', (1-2k)a^2+\frac{3}{2}c^2(2m-1), (3-6k)a^2+\frac{1}{2}c^2(2m-1)$
24	$\frac{2ak\sigma\sigma_m\left(\sqrt{m}\sigma_m+i\sqrt{m'}\right)\mathrm{cn}(\xi\|k)\mathrm{sn}(\eta\|m)\mathrm{sn}(\xi\|k)}{c\left(\sigma_m\sqrt{m'}-i\sqrt{m}\right)(\mathrm{cn}(\eta\|m)+1)\mathrm{dn}(\xi\|k)}$	$16k^2, \frac{3}{2}c^2(2m-1)-2a^2(k-2), \frac{1}{2}c^2(2m-1)-6a^2(k-2)$
25	$\frac{ak\sigma\sigma_k\sigma_m\left(\sqrt{m}\sigma_m+i\sqrt{m'}\right)\mathrm{cn}(\xi\|k)\mathrm{sn}(\eta\|m)}{c\left(\sigma_m\sqrt{m'}-i\sqrt{m}\right)(\mathrm{cn}(\eta\|m)+1)\left(\sqrt{k'}-\mathrm{dn}(\xi\|k)\right)}$	$k^2, \frac{3}{2}c^2(2m-1)-\frac{1}{2}a^2(k-2), \frac{1}{2}c^2(2m-1)-\frac{3}{2}a^2(k-2)$
26	$\frac{a\sigma\sigma_k\sigma_m\sqrt{k'}\left(\sqrt{m}\sigma_m+i\sqrt{m'}\right)\mathrm{dn}(\xi\|k)\mathrm{sn}(\eta\|m)}{c\left(i\sqrt{m'}\sigma_m+\sqrt{m}\right)(\mathrm{cn}(\eta\|m)+1)\left(\sqrt{k}\mathrm{sn}(\xi\|k)-1\right)}$	$k'^2, \frac{3}{2}c^2(2m-1)-\frac{1}{2}a^2(k+1), \frac{1}{2}c^2(2m-1)-\frac{3}{2}a^2(k+1)$
27	$\frac{ia\sigma\sigma_m(1-2\theta[0.5-k])\left(\sqrt{m}\sigma_m+i\sqrt{m'}\right)\mathrm{sn}(\eta\|m)\mathrm{sn}(\xi\|k)}{c\left(i\sqrt{m'}\sigma_m+\sqrt{m}\right)(\mathrm{cn}(\eta\|m)+1)(\mathrm{cn}(\xi\|k)+1)}$	$1, \left(k-\frac{1}{2}\right)a^2+\frac{3}{2}c^2(2m-1), \left(3k-\frac{3}{2}\right)a^2+\frac{1}{2}c^2(2m-1)$
28	$\frac{ia\sqrt{k}\sigma\mathrm{sn}(\xi\|k)\left(\sqrt{m}\sigma_m\mathrm{sn}(\eta\|m)^2+1\right)}{2c\sqrt{\sigma_m}\left(\sigma_m+\sqrt{m}\right)\sqrt[4]{m}\mathrm{sn}(\eta\|m)}$	$\frac{k}{4\sqrt{m}\sigma_m\left(\sigma_m+\sqrt{m}\right)^2}, (k+1)a^2+3c^2(m+1)+18c^2\sqrt{m}\sigma_m, 3(k+1)a^2+c^2(m+1)+6c^2\sqrt{m}\sigma_m$
29	$\frac{ia\sigma\sqrt{k'}\mathrm{sc}(\xi\|k)\left(\sqrt{m}\sigma_m\mathrm{sn}(\eta\|m)^2+1\right)}{2c\sqrt[4]{m}\sqrt{\sigma_m}\left(\sigma_m+\sqrt{m}\right)\mathrm{sn}(\eta\|m)}$	$\frac{k'}{4\sqrt{m}\sigma_m\left(\sigma_m+\sqrt{m}\right)^2}, (k-2)a^2+3c^2(m+1)+18c^2\sqrt{m}\sigma_m, 3(k-2)a^2+c^2(m+1)+6c^2\sqrt{m}\sigma_m$
30	$\frac{a\sigma\sqrt{kk'}\mathrm{sd}(\xi\|k)\left(\sqrt{m}\sigma_m\mathrm{sn}(\eta\|m)^2+1\right)}{2c\sqrt{\sigma_m}\left(\sigma_m+\sqrt{m}\right)\sqrt[4]{m}\mathrm{sn}(\eta\|m)}$	$-\frac{kk'}{4\sqrt{m}\sigma_m\left(\sigma_m+\sqrt{m}\right)^2}, (1-2k)a^2+3c^2(m+1)+18c^2\sqrt{m}\sigma_m, (3-6k)a^2+c^2(m+1)+6c^2\sqrt{m}\sigma_m$
31	$\frac{iak\sigma\mathrm{cn}(\xi\|k)\mathrm{sn}(\xi\|k)\left(\sqrt{m}\sigma_m\mathrm{sn}(\eta\|m)^2+1\right)}{2c\sqrt{\sigma_m}\left(\sigma_m+\sqrt{m}\right)\sqrt[4]{m}\mathrm{dn}(\xi\|k)\mathrm{sn}(\eta\|m)}$	$\frac{k^2}{4\sqrt{m}\sigma_m\left(\sigma_m+\sqrt{m}\right)^2}, -2(k-2)a^2+3c^2(m+1)+18c^2\sqrt{m}\sigma_m, -6(k-2)a^2+c^2(m+1)+6c^2\sqrt{m}\sigma_m$
32	$\frac{iak\sigma\mathrm{cn}(\xi\|k)\sigma_k\left(\sqrt{m}\sigma_m\mathrm{sn}(\eta\|m)^2+1\right)}{4c\sqrt[4]{m}\sqrt{\sigma_m}\left(\sigma_m+\sqrt{m}\right)\mathrm{sn}(\eta\|m)\left(\sqrt{k'}-\mathrm{dn}(\xi\|k)\right)}$	$\frac{k^2}{64\sqrt{m}\sigma_m\left(\sigma_m+\sqrt{m}\right)^2}, \frac{1}{2}(2-k)a^2+3c^2(m+1)+18c^2\sqrt{m}\sigma_m, -\frac{3}{2}(k-2)a^2+c^2(m+1)+6c^2\sqrt{m}\sigma_m$
33	$\frac{a\sigma\sigma_k\sqrt{k'}\mathrm{dn}(\xi\|k)\left(\sqrt{m}\sigma_m\mathrm{sn}(\eta\|m)^2+1\right)}{4c\sqrt[4]{m}\sqrt{\sigma_m}\left(\sigma_m+\sqrt{m}\right)\mathrm{sn}(\eta\|m)\left(\sqrt{k}\mathrm{sn}(\xi\|k)-1\right)}$	$\frac{k'^2}{64\sqrt{m}\sigma_m\left(\sigma_m+\sqrt{m}\right)^2}, \frac{1}{2}(-k-1)a^2+3c^2(m+1)+18c^2\sqrt{m}\sigma_m, -\frac{3}{2}(k+1)a^2+c^2(m+1)+6c^2\sqrt{m}\sigma_m$
34	$\frac{ia\sigma\left(1-2\theta[0.5-k]\mathrm{sn}(\xi\|k)\left(\sqrt{m}\sigma_m\mathrm{sn}(\eta\|m)^2+1\right)\right)}{4c\sqrt[4]{m}\sqrt{\sigma_m}\left(\sigma_m+\sqrt{m}\right)(\mathrm{cn}(\xi\|k)+1)\mathrm{sn}(\eta\|m)}$	$\frac{1}{64\sqrt{m}\sigma_m\left(\sigma_m+\sqrt{m}\right)^2}, \left(k-\frac{1}{2}\right)a^2+3c^2(m+1)+18c^2\sqrt{m}\sigma_m, \left(3k-\frac{3}{2}\right)a^2+c^2(m+1)+6c^2\sqrt{m}\sigma_m$
35	$\frac{ia\sigma\left(\sqrt{k}\sigma_k\mathrm{sn}(\xi\|k)^2+1\right)\left(\sqrt{m}\sigma_m\mathrm{sn}(\eta\|m)^2+1\right)}{2c\sqrt[4]{m}\sqrt{\sigma_m}\left(\sigma_m+\sqrt{m}\right)\mathrm{sn}(\eta\|m)\mathrm{sn}(\xi\|k)}$	$\frac{\sqrt{\frac{k}{m}}\sigma_k\left(\sigma_k+\sqrt{k}\right)^2}{\sigma_m\left(\sigma_m+\sqrt{m}\right)^2}, (k+1)a^2+6\sqrt{k}\sigma_k a^2+3c^2(m+1)+18c^2\sqrt{m}\sigma_m, 3(k+1)a^2+18\sqrt{k}\sigma_k a^2+c^2(m+1)+6c^2\sqrt{m}\sigma_m$
36	$\frac{ia\sigma\sqrt{k}\sqrt{\sigma_m}\left(m+\sigma_m\sqrt{-mm'}\right)}{2c\sqrt[4]{-mm'}}\mathrm{sn}(\xi\|k)\times\left(\mathrm{cn}(\eta\|m)-\frac{\sigma_m\sqrt{-\frac{m'}{m}}}{\mathrm{cn}(\eta\|m)}\right)$	$-\frac{k}{4\sigma_m\left(\sqrt{-m'}-\sqrt{m}\sigma_m\right)^2\sqrt{-mm'}}, (k+1)a^2+3c^2(1-2m)+18c^2\sigma_m\sqrt{-mm'}, 3(k+1)a^2+c^2(1-2m)+6c^2\sigma_m\sqrt{-mm'}$
37	$\frac{ia\sigma\sqrt{\sigma_m}\sqrt{k'}\left(m+\sigma_m\sqrt{-mm'}\right)}{2c\sqrt[4]{-mm'}}\mathrm{sc}(\xi\|k)\times\left(\mathrm{cn}(\eta\|m)-\frac{\sigma_m\sqrt{-\frac{m'}{m}}}{\mathrm{cn}(\eta\|m)}\right)$	$-\frac{k'}{4\sigma_m\left(\sqrt{-m'}-\sqrt{m}\sigma_m\right)^2\sqrt{-mm'}}, (k-2)a^2+3c^2(1-2m)+18c^2\sigma_m\sqrt{-mm'}, 3(k-2)a^2+c^2(1-2m)+6c^2\sigma_m\sqrt{-mm'}$

continue I

number 2 $\sqrt{\nu}\frac{\partial \tan^{-1}}{\partial x}$		$\frac{c^4}{a^4}, \frac{b}{a}, \frac{d}{c}$
38	$\frac{a\sigma\ \sqrt{\sigma_m}\ \sqrt{kk'}\ \left(m+\sigma_m\sqrt{-mm'}\right)}{2c\ \sqrt[4]{-mm'}}\mathrm{sd}(\xi\|k)$ $\times\frac{\left(\mathrm{cn}(\eta\|m)^2-\sigma_m\sqrt{-\frac{m'}{m}}\right)}{\mathrm{cn}(\eta\|m)}$	$\frac{kk'}{4\sigma_m\left(\sqrt{-m'}-\sqrt{m}\sigma_m\right)^2\sqrt{-mm'}}, (1-2k)a^2+3c^2(1-2m)+18c^2\sigma_m\sqrt{-mm'}, (3-6k)a^2+c^2(1-2m)+6c^2\sigma_m\sqrt{-mm'}$
39	$\frac{iak\sigma\ \sqrt{\sigma_m}\left(m+\sigma_m\sqrt{-mm'}\right)\mathrm{cn}(\xi\|k)\mathrm{sn}(\xi\|k)}{2c\ \sqrt[4]{-mm'}\mathrm{dn}(\xi\|k)}$ $\times\left(\mathrm{cn}(\eta\|m)-\frac{\sigma_m\sqrt{-\frac{m'}{m}}}{\mathrm{cn}(\eta\|m)}\right)$	$-\frac{k^2}{4\sigma_m\left(\sqrt{-m'}-\sqrt{m}\sigma_m\right)^2\sqrt{-mm'}}, -2(k-2)a^2+3c^2(1-2m)+18c^2\sigma_m\sqrt{-mm'}, -6(k-2)a^2+c^2(1-2m)+6c^2\sigma_m\sqrt{-mm'}$
40	$\frac{iak\sigma\ \sigma_k\ \sqrt{\sigma_m}\left(m+\sigma_m\sqrt{-mm'}\right)\mathrm{cn}(\xi\|k)}{4c\ \sqrt[4]{-mm'}\left(\sqrt{k'}-\mathrm{dn}(\xi\|k)\right)}$ $\times\left(\mathrm{cn}(\eta\|m)-\frac{\sigma_m\sqrt{-\frac{m'}{m}}}{\mathrm{cn}(\eta\|m)}\right)$	$-\frac{k^2}{64\sigma_m\left(\sqrt{-m'}-\sqrt{m}\sigma_m\right)^2\sqrt{-mm'}}, -\frac{1}{2}(k-2)a^2+3c^2(1-2m)+18c^2\sigma_m\sqrt{-mm'}, -\frac{3}{2}(k-2)a^2+c^2(1-2m)+6c^2\sigma_m\sqrt{-mm'}$
41	$\frac{a\sigma\ \sigma_k\ \sqrt{\sigma_m}\ \sqrt{k'}\left(m+\sigma_m\sqrt{-mm'}\right)\mathrm{dn}(\xi\|k)}{4c\ \sqrt[4]{-mm'}\left(\sqrt{k}\mathrm{sn}(\xi\|k)-1\right)}$ $\times\left(\mathrm{cn}(\eta\|m)-\frac{\sigma_m\sqrt{-\frac{m'}{m}}}{\mathrm{cn}(\eta\|m)}\right)$	$-\frac{k'^2}{64\sigma_m^3\left(\sqrt{-m'}-\sqrt{m}\sigma_m\right)^2\sqrt{-mm'}}, -\frac{1}{2}(k+1)a^2+3c^2(1-2m)+18c^2\sigma_m\sqrt{-mm'}, -\frac{3}{2}(k+1)a^2+c^2(1-2m)+6c^2\sigma_m\sqrt{-mm'}$
42	$\frac{ia\sigma\ \sqrt{\sigma_m}(1-2\theta[0.5-k])\left(m+\sigma_m\sqrt{-mm'}\right)\mathrm{sn}(\xi\|k)}{4c\ \sqrt[4]{-mm'}(\mathrm{cn}(\xi\|k)+1)}$ $\times\left(\mathrm{cn}(\eta\|m)-\frac{\sigma_m\sqrt{-\frac{m'}{m}}}{\mathrm{cn}(\eta\|m)}\right)$	$-\frac{1}{64\sigma_m\left(\sqrt{-m'}-\sqrt{m}\sigma_m\right)^2\sqrt{-mm'}}, \left(k-\frac{1}{2}\right)a^2+3c^2(1-2m)+18c^2\sigma_m\sqrt{-mm'}, \left(3k-\frac{3}{2}\right)a^2+c^2(1-2m)+6c^2\sigma_m\sqrt{-mm'}$
43	$\frac{ia\sigma\ \sqrt{\sigma_m}\left(m+\sigma_m\sqrt{-mm'}\right)\left(\sqrt{k}\sigma_k\mathrm{sn}(\xi\|k)^2+1\right)}{2c\ \sqrt[4]{-mm'}\mathrm{sn}(\xi\|k)}$ $\times\left(\mathrm{cn}(\eta\|m)-\frac{\sigma_m\sqrt{-\frac{m'}{m}}}{\mathrm{cn}(\eta\|m)}\right)$	$-\frac{k\sigma_k\left(\sigma_k+\sqrt{k}\right)^2}{\sigma_m\left(\sqrt{-m'}-\sqrt{m}\sigma_m\right)^2\sqrt{-kmm'}}, (k+1)a^2+6\sqrt{k}\sigma_k a^2+3c^2(1-2m)+18c^2\sigma_m\sqrt{-mm'}, 3(k+1)a^2+18\sqrt{k}\sigma_k a^2+c^2(1-2m)+6c^2\sigma_m\sqrt{-mm'}$
44	$\frac{ia\ \sigma\sigma_k\ \sqrt{k}\ \sqrt{\sigma_m}\left(m+\sigma_m\sqrt{-mm'}\right)}{2c\ \sqrt[4]{-mm'}}$ $\times\left(\mathrm{cn}(\xi\|k)-\frac{\sigma_k\sqrt{-\frac{k'}{k}}}{\mathrm{cn}(\xi\|k)}\right)\left(\mathrm{cn}(\eta\|m)-\frac{\sigma_m\sqrt{-\frac{m'}{m}}}{\mathrm{cn}(\eta\|m)}\right)$	$\frac{\sigma_k\left(\sqrt{-k'}-\sqrt{k}\sigma_k\right)^2\sqrt{-kk'}}{\sigma_m\left(\sqrt{-m'}-\sqrt{m}\sigma_m\right)^2\sqrt{-mm'}}, (1-2k)a^2+6\sigma_k\sqrt{-kk'}a^2+3c^2(1-2m)+18c^2\sigma_m\sqrt{-mm'}, (3-6k)a^2+18\sigma_k\sqrt{-kk'}a^2+c^2(1-2m)+6c^2\sigma_m\sqrt{-mm'}$
45	$\frac{ia\sigma\ \sqrt{k}\ \sqrt{\sigma_m}\mathrm{sn}(\xi\|k)\left(\sqrt{m'}-\mathrm{dn}(\eta\|m)^2\sigma_m\right)}{2c\left(\sigma_m\sqrt{m'}-1\right)\sqrt[4]{m'}\mathrm{dn}(\eta\|m)}$	$-\frac{k}{4\sigma_m\left(\sqrt{m'}-\sigma_m\right)^2\sqrt{m'}}, (k+1)a^2+3c^2(m-2)+18c^2\sigma_m\sqrt{m'}, 3(k+1)a^2+c^2(m-2)+6c^2\sigma_m\sqrt{m'}$
46	$\frac{ia\sigma\ \sqrt{\sigma_m}\ \sqrt{k'}\mathrm{sc}(\xi\|k)\left(\sqrt{m'}-\mathrm{dn}(\eta\|m)^2\sigma_m\right)}{2c\left(\sigma_m\sqrt{m'}-1\right)\sqrt[4]{m'}\mathrm{dn}(\eta\|m)}$	$-\frac{k'}{4\sigma_m\left(\sqrt{m'}-\sigma_m\right)^2\sqrt{m'}}, (k-2)a^2+3c^2(m-2)+18c^2\sigma_m\sqrt{m'}, 3(k-2)a^2+c^2(m-2)+6c^2\sigma_m\sqrt{m'}$
47	$\frac{a\sigma\ \sqrt{\sigma_m}\ \sqrt{kk'}\mathrm{sd}(\xi\|k)\left(\sqrt{m'}-\mathrm{dn}(\eta\|m)^2\sigma_m\right)}{2c\left(\sigma_m\sqrt{m'}-1\right)\sqrt[4]{m'}\mathrm{dn}(\eta\|m)}$	$\frac{kk'}{4\sigma_m\left(\sqrt{m'}-\sigma_m\right)^2\sqrt{m'}}, (1-2k)a^2+3c^2(m-2)+18c^2\sigma_m\sqrt{m'}, (3-6k)a^2+c^2(m-2)+6c^2\sigma_m\sqrt{m'}$
48	$\frac{iak\sigma\ \sqrt{\sigma_m}\mathrm{cn}(\xi\|k)\mathrm{sn}(\xi\|k)\left(\sqrt{m'}-\mathrm{dn}(\eta\|m)^2\sigma_m\right)}{2c\left(\sigma_m\sqrt{m'}-1\right)\sqrt[4]{m'}\mathrm{dn}(\eta\|m)\mathrm{dn}(\xi\|k)}$	$-\frac{k^2}{4\sigma_m\left(\sqrt{m'}-\sigma_m\right)^2\sqrt{m'}}, -2(k-2)a^2+3c^2(m-2)+18c^2\sigma_m\sqrt{m'}, -6(k-2)a^2+c^2(m-2)+6c^2\sigma_m\sqrt{m'}$
49	$\frac{iak\sigma\ \sigma_k\ \sqrt{\sigma_m}\mathrm{cn}(\xi\|k)\left(\mathrm{dn}(\eta\|m)^2\sigma_m-\sqrt{m'}\right)}{4c\left(\sigma_m\sqrt{m'}-1\right)\sqrt[4]{m'}\mathrm{dn}(\eta\|m)\left(\sqrt{k'}-\mathrm{dn}(\xi\|k)\right)}$	$-\frac{k^2}{64\sigma_m\left(\sqrt{m'}-\sigma_m\right)^2\sqrt{m'}}, -\frac{1}{2}(k-2)a^2+3c^2(m-2)+18c^2\sigma_m\sqrt{m'}, -\frac{3}{2}(k-2)a^2+c^2(m-2)+6c^2\sigma_m\sqrt{m'}$
50	$\frac{a\sigma\ \sigma_k\ \sqrt{\sigma_m}\ \sqrt{k'}\mathrm{dn}(\xi\|k)\left(\mathrm{dn}(\eta\|m)^2\sigma_m-\sqrt{m'}\right)}{4c\left(\sigma_m\sqrt{m'}-1\right)\sqrt[4]{m'}\mathrm{dn}(\eta\|m)\left(\sqrt{k}\mathrm{sn}(\xi\|k)-1\right)}$	$-\frac{k'^2}{64\sigma_m\left(\sqrt{m'}-\sigma_m\right)^2\sqrt{m'}}, -\frac{1}{2}(k+1)a^2+3c^2(m-2)+18c^2\sigma_m\sqrt{m'}, -\frac{3}{2}(k+1)a^2+c^2(m-2)+6c^2\sigma_m\sqrt{m'}$
51	$\frac{a\sigma i(1-2\theta[0.5-k])\ \sqrt{\sigma_m}\mathrm{sn}(\xi\|k)\left(\sqrt{m'}-\mathrm{dn}(\eta\|m)^2\sigma_m\right)}{4c\left(\sigma_m\sqrt{m'}-1\right)\sqrt[4]{m'}(\mathrm{cn}(\xi\|k)+1)\mathrm{dn}(\eta\|m)}$	$-\frac{1}{64\sigma_m\left(\sqrt{m'}-\sigma_m\right)^2\sqrt{m'}}, \left(k-\frac{1}{2}\right)a^2+3c^2(m-2)+18c^2\sigma_m\sqrt{m'}, \left(3k-\frac{3}{2}\right)a^2+c^2(m-2)+6c^2\sigma_m\sqrt{m'}$
52	$\frac{ia\sigma\ \sqrt{\sigma_m}\left(\sqrt{k}\sigma_k\mathrm{sn}(\xi\|k)^2+1\right)\left(\sqrt{m'}-\mathrm{dn}(\eta\|m)^2\sigma_m\right)}{2c\left(\sigma_m\sqrt{m'}-1\right)\sqrt[4]{m'}\mathrm{dn}(\eta\|m)\mathrm{sn}(\xi\|k)}$	$-\frac{\sqrt{k-km}\sigma_k\left(\sigma_k+\sqrt{k}\right)^2}{\sigma_m\left(\sqrt{m'}-\sigma_m\right)^2m'}, (k+1)a^2+6\sqrt{k}\sigma_k a^2+3c^2(m-2)+18c^2\sigma_m\sqrt{m'}, 3(k+1)a^2+18\sqrt{k}\sigma_k a^2+c^2(m-2)+6c^2\sigma_m\sqrt{m'}$
53	$\frac{a\sqrt{k}\sigma\sigma_k\ \sqrt{\sigma_m}\left(\mathrm{dn}(\eta\|m)^2\sigma_m-\sqrt{m'}\right)}{2c\left(\sigma_m\sqrt{m'}-1\right)\sqrt[4]{m'}\mathrm{dn}(\eta\|m)}$ $\times\left(\mathrm{cn}(\xi\|k)-\frac{\sigma_k\sqrt{-\frac{k'}{k}}}{\mathrm{cn}(\xi\|k)}\right)$	$\frac{\sigma_k\left(\sqrt{-k'}-\sqrt{k}\sigma_k\right)^2\sqrt{-\frac{kk'}{m'}}}{\sigma_m\left(\sqrt{m'}-\sigma_m\right)^2}, (1-2k)a^2+6\sigma_k\sqrt{-kk'}a^2+3c^2(m-2)+18c^2\sigma_m\sqrt{m'}, (3-6k)a^2+18\sigma_k\sqrt{-kk'}a^2+c^2(m-2)+6c^2\sigma_m\sqrt{m'}$
54	$\frac{a\sigma\sigma_k\ \sqrt{\sigma_m}\left(\sqrt{k'}-\mathrm{dn}(\xi\|k)^2\sigma_k\right)\left(\sqrt{m'}-\mathrm{dn}(\eta\|m)^2\sigma_m\right)}{2c\left(\sigma_m\sqrt{m'}-1\right)\sqrt[4]{m'}\mathrm{dn}(\eta\|m)\mathrm{dn}(\xi\|k)}$	$\frac{\sigma_k\left(\sqrt{k'}-\sigma_k\right)^2\sqrt{\frac{k'}{m'}}}{\sigma_m\left(\sqrt{m'}-\sigma_m\right)^2}, (k-2)a^2+6\sigma_k\sqrt{k'}a^2+3c^2(m-2)+18c^2\sigma_m\sqrt{m'}, 3(k-2)a^2+18\sigma_k\sqrt{k'}a^2+c^2(m-2)+6c^2\sigma_m\sqrt{m'}$
55	$\frac{2a\sqrt{k}\sigma\mathrm{sn}(\xi\|k)\left(\sigma_m^{3/2}+\sqrt[4]{m}\mathrm{sn}(\eta\|m)\right)}{c\left(\sqrt{m}-\sigma_m\right)\sigma_m\left(\sigma_m^{3/2}-\sqrt[4]{m}\mathrm{sn}(\eta\|m)\right)}$	$\frac{16k}{\left(\sqrt{m}\sigma_m-1\right)^4}, (k+1)a^2-\frac{3}{2}c^2(m+1)-9c^2\sqrt{m}\sigma_m, 3(k+1)a^2-\frac{c^2}{2}(m+1)-3c^2\sqrt{m}\sigma_m$

continue I

number $2\sqrt{\nu}\frac{\partial\tan^{-1}}{\partial x}$		$\frac{c^4}{a^4}, \frac{b}{a}, \frac{d}{c}$
56	$\frac{2a\sigma\sqrt{k'}\,\mathrm{sc}(\xi\|k)\left(\sigma_m^{3/2}+\sqrt[4]{m}\,\mathrm{sn}(\eta\|m)\right)}{c\left(\sqrt{m}-\sigma_m\right)\sigma_m\left(\sigma_m^{3/2}-\sqrt[4]{m}\,\mathrm{sn}(\eta\|m)\right)}$	$\frac{16k'}{\left(\sqrt{m}\sigma_m-1\right)^4}, (k-2)a^2-\frac{3}{2}c^2(m+1)-9c^2\sqrt{m}\sigma_m, 3(k-2)a^2-\frac{1}{2}c^2(m+1)-3c^2\sqrt{m}\sigma_m$
57	$\frac{2ia\sigma\sqrt{kk'}\,\mathrm{sd}(\xi\|k)\left(\sigma_m^{3/2}+\sqrt[4]{m}\,\mathrm{sn}(\eta\|m)\right)}{c\left(\sqrt{m}-\sigma_m\right)\sigma_m\left(\sigma_m^{3/2}-\sqrt[4]{m}\,\mathrm{sn}(\eta\|m)\right)}$	$-\frac{16kk'}{\left(\sqrt{m}\sigma_m-1\right)^4}, (1-2k)a^2-\frac{3}{2}c^2(m+1)-9c^2\sqrt{m}\sigma_m, (3-6k)a^2-\frac{1}{2}c^2(m+1)-3c^2\sqrt{m}\sigma_m$
58	$\frac{2ak\sigma\,\mathrm{cn}(\xi\|k)\mathrm{sn}(\xi\|k)\left(\sigma_m^{3/2}+\sqrt[4]{m}\,\mathrm{sn}(\eta\|m)\right)}{c\left(\sqrt{m}-\sigma_m\right)\sigma_m\,\mathrm{dn}(\xi\|k)\left(\sigma_m^{3/2}-\sqrt[4]{m}\,\mathrm{sn}(\eta\|m)\right)}$	$\frac{16k^2}{\left(\sqrt{m}\sigma_m-1\right)^4}, \left(-2(k-2)a^2-\frac{3}{2}c^2(m+1)\right)-9c^2\sqrt{m}\sigma_m, -6(k-2)a^2-\frac{1}{2}c^2(m+1)-3c^2\sqrt{m}\sigma_m$
59	$\frac{ak\sigma\sigma_k\,\mathrm{cn}(\xi\|k)\left(\sigma_m^{3/2}+\sqrt[4]{m}\,\mathrm{sn}(\eta\|m)\right)}{c\left(\sqrt{m}-\sigma_m\right)\sigma_m\left(\sigma_m^{3/2}-\sqrt[4]{m}\,\mathrm{sn}(\eta\|m)\right)\left(\sqrt{k'}-\mathrm{dn}(\xi\|k)\right)}$	$\frac{k^2}{\left(\sqrt{m}\sigma_m-1\right)^4}, \frac{1}{2}(2-k)a^2-\frac{3}{2}c^2(m+1)-9c^2\sqrt{m}\sigma_m, -\frac{3}{2}(k-2)a^2-\frac{c^2}{2}(m+1)-3c^2\sqrt{m}\sigma_m$
60	$\frac{ia\sigma\sigma_k\sqrt{k'}\,\mathrm{dn}(\xi\|k)\left(\sigma_m^{3/2}+\sqrt[4]{m}\,\mathrm{sn}(\eta\|m)\right)}{c\left(\sqrt{m}-\sigma_m\right)\sigma_m\left(\sqrt{k}\,\mathrm{sn}(\xi\|k)-1\right)\left(\sigma_m^{3/2}-\sqrt[4]{m}\,\mathrm{sn}(\eta\|m)\right)}$	$\frac{k'^2}{\left(\sqrt{m}\sigma_m-1\right)^4}, \frac{1}{2}(-k-1)a^2-\frac{3}{2}c^2(m+1)-9c^2\sqrt{m}\sigma_m, -\frac{3}{2}(k+1)a^2-\frac{c^2}{2}(m+1)-3c^2\sqrt{m}\sigma_m$
61	$\frac{a\sigma(1-2\theta[0.5-k])\mathrm{sn}(\xi\|k)\left(\sigma_m^{3/2}+\sqrt[4]{m}\,\mathrm{sn}(\eta\|m)\right)}{c\left(\sqrt{m}-\sigma_m\right)\sigma_m(\mathrm{cn}(\xi\|k)+1)\left(\sigma_m^{3/2}-\sqrt[4]{m}\,\mathrm{sn}(\eta\|m)\right)}$	$\frac{1}{\left(\sqrt{m}\sigma_m-1\right)^4}, \frac{1}{2}(2k-1)a^2-\frac{3}{2}c^2(m+1)-9c^2\sqrt{m}\sigma_m, \frac{1}{2}(6k-3)a^2-\frac{c^2}{2}(m+1)-3c^2\sqrt{m}\sigma_m$
62	$\frac{2a\sigma\left(\sqrt{k}\sigma_k\,\mathrm{sn}(\xi\|k)^2+1\right)\left(\sigma_m^{3/2}+\sqrt[4]{m}\,\mathrm{sn}(\eta\|m)\right)}{c\sigma_m\left(\sqrt{m}-\sigma_m\right)\mathrm{sn}(\xi\|k)\left(\sigma_m^{3/2}-\sqrt[4]{m}\,\mathrm{sn}(\eta\|m)\right)}$	$\frac{64\sqrt{k}\sigma_k\left(\sigma_k+\sqrt{k}\right)^2}{\left(\sqrt{m}\sigma_m-1\right)^4}, (k+1)a^2+6\sqrt{k}\sigma_k a^2-\frac{3}{2}c^2(m+1)-9c^2\sqrt{m}\sigma_m, 3(k+1)a^2+18\sqrt{k}\sigma_k a^2-\frac{c^2}{2}(m+1)-3c^2\sqrt{m}\sigma_m$
63	$\frac{2ia\sqrt{k}\sigma\sigma_k\left(\sigma_m^{3/2}+\sqrt[4]{m}\,\mathrm{sn}(\eta\|m)\right)\left(\mathrm{cn}(\xi\|k)-\frac{\sigma_k\sqrt{-\frac{k'}{k}}}{\mathrm{cn}(\xi\|k)}\right)}{c\left(\sqrt{m}-\sigma_m\right)\sigma_m\left(\sigma_m^{3/2}-\sqrt[4]{m}\,\mathrm{sn}(\eta\|m)\right)}$	$-\frac{64\sigma_k\left(\sqrt{-k'}-\sqrt{k}\sigma_k\right)^2\sqrt{-kk'}}{\left(\sqrt{m}\sigma_m-1\right)^4}, (1-2k)a^2+6\sigma_k\sqrt{-kk'}a^2-\frac{3}{2}c^2(m+1)-9c^2\sqrt{m}\sigma_m, 3(1-2k)a^2+18\sigma_k\sqrt{-kk'}a^2-\frac{c^2}{2}(m+1)-3c^2\sqrt{m}\sigma_m$
64	$\frac{2ia\sigma\sigma_k\left(\sigma_m^{3/2}+\sqrt[4]{m}\,\mathrm{sn}(\eta\|m)\right)\left(\sqrt{k'}-\mathrm{dn}(\xi\|k)^2\sigma_k\right)}{c\sigma_m\left(\sqrt{m}-\sigma_m\right)\mathrm{dn}(\xi\|k)\left(\sigma_m^{3/2}-\sqrt[4]{m}\,\mathrm{sn}(\eta\|m)\right)}$	$-\frac{64\sigma_k\left(\sqrt{k'}-\sigma_k\right)^2\sqrt{k'}}{\left(\sqrt{m}\sigma_m-1\right)^4}, (k-2)a^2+6\sigma_k\sqrt{k'}a^2-\frac{3}{2}c^2(m+1)-9c^2\sqrt{m}\sigma_m, 3(k-2)a^2+18\sigma_k\sqrt{k'}a^2-\frac{1}{2}c^2(m+1)-3c^2\sqrt{m}\sigma_m$
65	$\frac{a\sigma i\left(\sqrt{k}-\sigma_k\right)\left(\sigma_k^{3/2}+\sqrt[4]{k}\,\mathrm{sn}(\xi\|k)\right)\left(\sigma_m^{3/2}+\sqrt[4]{m}\,\mathrm{sn}(\eta\|m)\right)}{c\sigma_m\left(\sqrt{m}-\sigma_m\right)\left(\sigma_k^{3/2}-\sqrt[4]{k}\,\mathrm{sn}(\xi\|k)\right)\left(\sigma_m^{3/2}-\sqrt[4]{m}\,\mathrm{sn}(\eta\|m)\right)}$	$\frac{\left(\sqrt{k}\sigma_k-1\right)^4}{\left(\sqrt{m}\sigma_m-1\right)^4}, \frac{1}{2}(-k-1)a^2-3\sqrt{k}\sigma_k a^2-\frac{3}{2}c^2(m+1)-9c^2\sqrt{m}\sigma_m, -\frac{3}{2}(k+1)a^2-9\sqrt{k}\sigma_k a^2-\frac{c^2}{2}(m+1)-3c^2\sqrt{m}\sigma_m$
66	$\frac{2a\sqrt{k}\sigma\,\mathrm{sn}(\xi\|k)\left(\sqrt[4]{m'}(-\sigma_m)^{3/2}+\sqrt[4]{-m}\,\mathrm{cn}(\eta\|m)\right)}{c\left(\sqrt{m'}\sigma_m+\sqrt{-m}\right)\left(\sqrt[4]{-m}\,\mathrm{cn}(\eta\|m)-(-\sigma_m)^{3/2}\sqrt[4]{m'}\right)}$	$\frac{16k}{\left(\sqrt{m'}\sigma_m+\sqrt{-m}\right)^4}, (k+1)a^2+\frac{3}{2}c^2(2m-1)+9c^2\sigma_m\sqrt{-mm'}, 3(k+1)a^2+\frac{1}{2}c^2(2m-1)+3c^2\sigma_m\sqrt{-mm'}$
67	$\frac{2a\sigma\sqrt{k'}\,\mathrm{sc}(\xi\|k)\left(\sqrt[4]{m'}(-\sigma_m)^{3/2}+\sqrt[4]{-m}\,\mathrm{cn}(\eta\|m)\right)}{c\left(\sqrt{m'}\sigma_m+\sqrt{-m}\right)\left(\sqrt[4]{-m}\,\mathrm{cn}(\eta\|m)-(-\sigma_m)^{3/2}\sqrt[4]{m'}\right)}$	$\frac{16k'}{\left(\sqrt{m'}\sigma_m+\sqrt{-m}\right)^4}, (k-2)a^2+\frac{3}{2}c^2(2m-1)+9c^2\sigma_m\sqrt{-mm'}, 3(k-2)a^2+\frac{1}{2}c^2(2m-1)+3c^2\sigma_m\sqrt{-mm'}$
68	$\frac{2ia\sigma\sqrt{kk'}\,\mathrm{sd}(\xi\|k)\left(\sqrt[4]{m'}(-\sigma_m)^{3/2}+\sqrt[4]{-m}\,\mathrm{cn}(\eta\|m)\right)}{c\left(\sqrt{m'}\sigma_m+\sqrt{-m}\right)\left(\sqrt[4]{-m}\,\mathrm{cn}(\eta\|m)-(-\sigma_m)^{3/2}\sqrt[4]{m'}\right)}$	$-\frac{16kk'}{\left(\sqrt{m'}\sigma_m+\sqrt{-m}\right)^4}, (1-2k)a^2+\frac{3}{2}c^2(2m-1)+9c^2\sigma_m\sqrt{-mm'}, (3-6k)a^2+\frac{1}{2}c^2(2m-1)+3c^2\sigma_m\sqrt{-mm'}$
69	$\frac{2ak\sigma\,\mathrm{cn}(\xi\|k)\mathrm{sn}(\xi\|k)}{c\left(\sqrt{m'}\sigma_m+\sqrt{-m}\right)\mathrm{dn}(\xi\|k)} \times \frac{\left(\sqrt[4]{m'}(-\sigma_m)^{3/2}+\sqrt[4]{-m}\,\mathrm{cn}(\eta\|m)\right)}{\left(\sqrt[4]{-m}\,\mathrm{cn}(\eta\|m)-(-\sigma_m)^{3/2}\sqrt[4]{m'}\right)}$	$\frac{16k^2}{\left(\sqrt{m'}\sigma_m+\sqrt{-m}\right)^4}, -2(k-2)a^2+\frac{3}{2}c^2(2m-1)+9c^2\sigma_m\sqrt{-mm'}, -6(k-2)a^2+\frac{1}{2}c^2(2m-1)+3c^2\sigma_m\sqrt{-mm'}$
70	$\frac{ak\sigma\sigma_k\,\mathrm{cn}(\xi\|k)}{c\left(\sqrt{m'}\sigma_m+\sqrt{-m}\right)\left(\sqrt{k'}-\mathrm{dn}(\xi\|k)\right)} \times \frac{\left(\sqrt[4]{m'}(-\sigma_m)^{3/2}+\sqrt[4]{-m}\,\mathrm{cn}(\eta\|m)\right)}{\left(\sqrt[4]{-m}\,\mathrm{cn}(\eta\|m)-(-\sigma_m)^{3/2}\sqrt[4]{m'}\right)}$	$\frac{k^2}{\left(\sqrt{m'}\sigma_m+\sqrt{-m}\right)^4}, -\frac{1}{2}(k-2)a^2+\frac{3}{2}c^2(2m-1)+9c^2\sigma_m\sqrt{-mm'}, -\frac{3}{2}(k-2)a^2+\frac{c^2}{2}(2m-1)+3c^2\sigma_m\sqrt{-mm'}$
71	$\frac{ia\sigma\sigma_k\sqrt{k'}\,\mathrm{dn}(\xi\|k)}{c\left(\sqrt{m'}\sigma_m+\sqrt{-m}\right)\left(\sqrt{k}\,\mathrm{sn}(\xi\|k)-1\right)} \times \frac{\left(\sqrt[4]{m'}(-\sigma_m)^{3/2}+\sqrt[4]{-m}\,\mathrm{cn}(\eta\|m)\right)}{\left(\sqrt[4]{-m}\,\mathrm{cn}(\eta\|m)-(-\sigma_m)^{3/2}\sqrt[4]{m'}\right)}$	$\frac{k'^2}{\left(\sqrt{m'}\sigma_m+\sqrt{-m}\right)^4}, -\frac{1}{2}(k+1)a^2+\frac{3}{2}c^2(2m-1)+9c^2\sigma_m\sqrt{-mm'}, -\frac{3}{2}(k+1)a^2+\frac{c^2}{2}(2m-1)+3c^2\sigma_m\sqrt{-mm'}$
72	$\frac{a\sigma(1-2\theta[0.5-k])\mathrm{sn}(\xi\|k)}{c\left(\sqrt{m'}\sigma_m+\sqrt{-m}\right)(\mathrm{cn}(\xi\|k)+1)} \times \frac{\left(\sqrt[4]{m'}(-\sigma_m)^{3/2}+\sqrt[4]{-m}\,\mathrm{cn}(\eta\|m)\right)}{\left(\sqrt[4]{-m}\,\mathrm{cn}(\eta\|m)-(-\sigma_m)^{3/2}\sqrt[4]{m'}\right)}$	$\frac{1}{\left(\sqrt{m'}\sigma_m+\sqrt{-m}\right)^4}, \left(k-\frac{1}{2}\right)a^2+\frac{3}{2}c^2(2m-1)+9c^2\sigma_m\sqrt{-mm'}, \left(3k-\frac{3}{2}\right)a^2+\frac{1}{2}c^2(2m-1)+3c^2\sigma_m\sqrt{-mm'}$
73	$\frac{2a\sigma\left(\sqrt{k}\sigma_k\,\mathrm{sn}(\xi\|k)^2+1\right)}{c\left(\sqrt{m'}\sigma_m+\sqrt{-m}\right)\mathrm{sn}(\xi\|k)} \times \frac{\left(\sqrt[4]{m'}(-\sigma_m)^{3/2}+\sqrt[4]{-m}\,\mathrm{cn}(\eta\|m)\right)}{\left(\sqrt[4]{-m}\,\mathrm{cn}(\eta\|m)-(-\sigma_m)^{3/2}\sqrt[4]{m'}\right)}$	$\frac{64\sqrt{k}\sigma_k\left(\sigma_k+\sqrt{k}\right)^2}{\left(\sqrt{m'}\sigma_m+\sqrt{-m}\right)^4}, (k+1)a^2+6\sqrt{k}\sigma_k a^2+\frac{3}{2}c^2(2m-1)+9c^2\sigma_m\sqrt{-mm'}, 3(k+1)a^2+18\sqrt{k}\sigma_k a^2+\frac{1}{2}c^2(2m-1)+3c^2\sigma_m\sqrt{-mm'}$

continue I

number 2 $\sqrt{\nu}\frac{\partial\tan^{-1}}{\partial x}$	$\frac{c^4}{a^4}, \frac{b}{a}, \frac{d}{c}$
74 $\frac{2ia\sqrt{k}\sigma\sigma_k\left(\mathrm{cn}(\xi\|k)-\frac{\sigma_k\sqrt{-\frac{k'}{k}}}{\mathrm{cn}(\xi\|k)}\right)}{c\left(\sqrt{m'}\sigma_m+\sqrt{-m}\right)}$ $\times\frac{\left(\sqrt[4]{m'}(-\sigma_m)^{3/2}+\sqrt[4]{-m}\mathrm{cn}(\eta\|m)\right)}{\left(\sqrt[4]{-m}\mathrm{cn}(\eta\|m)-(-\sigma_m)^{3/2}\sqrt[4]{m'}\right)}$	$-\frac{64\sigma_k\left(\sqrt{-k'}-\sqrt{k}\sigma_k\right)^2\sqrt{-kk'}}{\left(\sqrt{m'}\sigma_m+\sqrt{-m}\right)^4}, (1-2k)a^2+6\sigma_k\sqrt{-kk'}a^2+\frac{3}{2}c^2(2m-1)+9c^2\sigma_m\sqrt{-mm'}, (3-6k)a^2+18\sigma_k\sqrt{-kk'}a^2+\frac{1}{2}c^2(2m-1)+3c^2\sigma_m\sqrt{-mm'}$
75 $\frac{2ia\sigma\sigma_k\left(\mathrm{dn}(\xi\|k)^2\sigma_k-\sqrt{k'}\right)}{c\left(\sqrt{m'}\sigma_m+\sqrt{-m}\right)\mathrm{dn}(\xi\|k)}$ $\times\frac{\left(\sqrt[4]{m'}(-\sigma_m)^{3/2}+\sqrt[4]{-m}\mathrm{cn}(\eta\|m)\right)}{\left(\sqrt[4]{-m}\mathrm{cn}(\eta\|m)-(-\sigma_m)^{3/2}\sqrt[4]{m'}\right)}$	$-\frac{64\sigma_k\left(\sqrt{k'}-\sigma_k\right)^2\sqrt{k'}}{\left(\sqrt{m'}\sigma_m+\sqrt{-m}\right)^4}, (k-2)a^2+6\sigma_k\sqrt{k'}a^2+\frac{3}{2}c^2(2m-1)+9c^2\sigma_m\sqrt{-mm'}, 3(k-2)a^2+18\sigma_k\sqrt{k'}a^2+\frac{1}{2}c^2(2m-1)+3c^2\sigma_m\sqrt{-mm'}$
76 $\frac{a\sigma\left(\sqrt{k}-\sigma_k\right)\left(\sigma_k^{3/2}+\sqrt[4]{k}\mathrm{sn}(\xi\|k)\right)}{c\left(\sqrt{m'}\sigma_m+\sqrt{-m}\right)\left(\sigma_k^{3/2}-\sqrt[4]{k}\mathrm{sn}(\xi\|k)\right)}$ $\times\frac{\left(\sqrt[4]{m'}(-\sigma_m)^{3/2}+\sqrt[4]{-m}\mathrm{cn}(\eta\|m)\right)}{\left(\sqrt[4]{-m}\mathrm{cn}(\eta\|m)-(-\sigma_m)^{3/2}\sqrt[4]{m'}\right)}$	$\frac{\left(\sqrt{k}\sigma_k-1\right)^4}{\left(\sqrt{m'}\sigma_m+\sqrt{-m}\right)^4}, \frac{1}{2}(-k-1)a^2-3\sqrt{k}\sigma_k a^2+\frac{3}{2}c^2(2m-1)+9c^2\sigma_m\sqrt{-mm'}, -\frac{3}{2}(k+1)a^2-9\sqrt{k}\sigma_k a^2+\frac{c^2}{2}(2m-1)+3c^2\sigma_m\sqrt{-mm'}$
77 $\frac{ia\sigma\left(\sqrt{k'}\sigma_k+\sqrt{-k}\right)\left(\sqrt[4]{k'}(-\sigma_k)^{3/2}+\sqrt[4]{-k}\mathrm{cn}(\xi\|k)\right)}{c\left(\sqrt{m'}\sigma_m+\sqrt{-m}\right)\left(\sqrt[4]{-k}\mathrm{cn}(\xi\|k)-(-\sigma_k)^{3/2}\sqrt[4]{k'}\right)}$ $\times\frac{\left(\sqrt[4]{m'}(-\sigma_m)^{3/2}+\sqrt[4]{-m}\mathrm{cn}(\eta\|m)\right)}{\left(\sqrt[4]{-m}\mathrm{cn}(\eta\|m)-(-\sigma_m)^{3/2}\sqrt[4]{m'}\right)}$	$\frac{\left(\sqrt{k'}\sigma_k+\sqrt{-k}\right)^4}{\left(\sqrt{m'}\sigma_m+\sqrt{-m}\right)^4}, \left(k-\frac{1}{2}\right)a^2+3\sigma_k\sqrt{-kk'}a^2+\frac{3}{2}c^2(2m-1)+9c^2\sigma_m\sqrt{-mm'}, 9\sigma_k\sqrt{-kk'}a^2+\frac{1}{2}(6k-3)a^2+\frac{c^2}{2}(2m-1)+3c^2\sigma_m\sqrt{-mm'}$
78 $\frac{2ia\sigma_m\sqrt{k}\sigma\mathrm{sn}(\xi\|k)\left(\sqrt[4]{m'}\sigma_m^{3/2}+\mathrm{dn}(\eta\|m)\right)}{c\left(\sigma_m\sqrt{m'}-1\right)\left(\mathrm{dn}(\eta\|m)-\sigma_m^{3/2}\sqrt[4]{m'}\right)}$	$\frac{16k}{\left(\sigma_m\sqrt{m'}-1\right)^4}, (k+1)a^2-\frac{3}{2}c^2(m-2)+9c^2\sigma_m\sqrt{m'}, 3(k+1)a^2-\frac{1}{2}c^2(m-2)+3c^2\sigma_m\sqrt{m'}$
79 $\frac{2ia\sigma\,\sigma_m\sqrt{k'}\mathrm{sc}(\xi\|k)\left(\sqrt[4]{m'}\sigma_m^{3/2}+\mathrm{dn}(\eta\|m)\right)}{c\left(\sigma_m\sqrt{m'}-1\right)\left(\mathrm{dn}(\eta\|m)-\sigma_m^{3/2}\sqrt[4]{m'}\right)}$	$\frac{16k'}{\left(\sigma_m\sqrt{m'}-1\right)^4}, (k-2)a^2-\frac{3}{2}c^2(m-2)+9c^2\sigma_m\sqrt{m'}, 3(k-2)a^2-\frac{1}{2}c^2(m-2)+3c^2\sigma_m\sqrt{m'}$
80 $\frac{2a\sigma\,\sigma_m\sqrt{kk'}\mathrm{sd}(\xi\|k)\left(\sqrt[4]{m'}\sigma_m^{3/2}+\mathrm{dn}(\eta\|m)\right)}{c\left(\sigma_m\sqrt{m'}-1\right)\left(\mathrm{dn}(\eta\|m)-\sigma_m^{3/2}\sqrt[4]{m'}\right)}$	$-\frac{16kk'}{\left(\sigma_m\sqrt{m'}-1\right)^4}, (1-2k)a^2-\frac{3}{2}c^2(m-2)+9c^2\sigma_m\sqrt{m'}, (3-6k)a^2-\frac{1}{2}c^2(m-2)+3c^2\sigma_m\sqrt{m'}$
81 $\frac{2iak\sigma\sigma_m\,\mathrm{cn}(\xi\|k)\mathrm{sn}(\xi\|k)\left(\sqrt[4]{m'}\sigma_m^{3/2}+\mathrm{dn}(\eta\|m)\right)}{c\left(\sigma_m\sqrt{m'}-1\right)\mathrm{dn}(\xi\|k)\left(\mathrm{dn}(\eta\|m)-\sigma_m^{3/2}\sqrt[4]{m'}\right)}$	$\frac{16k^2}{\left(\sigma_m\sqrt{m'}-1\right)^4}, -2(k-2)a^2-\frac{3}{2}c^2(m-2)+9c^2\sigma_m\sqrt{m'}, -6(k-2)a^2-\frac{1}{2}c^2(m-2)+3c^2\sigma_m\sqrt{m'}$
82 $\frac{iak\sigma\,\sigma_k\sigma_m\mathrm{cn}(\xi\|k)\left(\sigma_m^{3/2}\sqrt[4]{m'}+\mathrm{dn}(\eta\|m)\right)}{c\left(\sigma_m\sqrt{m'}-1\right)\left(\sqrt{k'}-\mathrm{dn}(\xi\|k)\right)\left(\mathrm{dn}(\eta\|m)-\sigma_m^{3/2}\sqrt[4]{m'}\right)}$	$\frac{k^2}{\left(\sigma_m\sqrt{m'}-1\right)^4}, -\frac{1}{2}(k-2)a^2-\frac{3}{2}c^2(m-2)+9c^2\sigma_m\sqrt{m'}, -\frac{3}{2}(k-2)a^2-\frac{c^2}{2}(m-2)+3c^2\sigma_m\sqrt{m'}$
83 $\frac{a\sigma\,\sigma_k\sigma_m\sqrt{k'}\mathrm{dn}(\xi\|k)\left(\sigma_m^{3/2}\sqrt[4]{m'}+\mathrm{dn}(\eta\|m)\right)}{c\left(\sigma_m\sqrt{m'}-1\right)\left(\sqrt{k}\mathrm{sn}(\xi\|k)-1\right)\left(\mathrm{dn}(\eta\|m)-\sigma_m^{3/2}\sqrt[4]{m'}\right)}$	$\frac{k'^2}{\left(\sigma_m\sqrt{m'}-1\right)^4}, -\frac{1}{2}(k+1)a^2-\frac{3}{2}c^2(m-2)+9c^2\sigma_m\sqrt{m'}, -\frac{3}{2}(k+1)a^2-\frac{c^2}{2}(m-2)+3c^2\sigma_m\sqrt{m'}$
84 $\frac{a\sigma\,\sigma_m(1-2\theta[0.5-k])\mathrm{sn}(\xi\|k)}{c\left(\sigma_m^{3/2}\sqrt[4]{m'}-1\right)\left(\sqrt[4]{m'}\sigma_m^{3/2}+1\right)(\mathrm{cn}(\xi\|k)+1)}$ $\times\frac{\left(\sigma_m^{3/2}\sqrt[4]{m'}+\mathrm{dn}(\eta\|m)\right)}{\left(\sigma_m^{3/2}\sqrt[4]{m'}-\mathrm{dn}(\eta\|m)\right)}$	$\frac{1}{\left(\sigma_m\sqrt{m'}-1\right)^4}, \left(k-\frac{1}{2}\right)a^2-\frac{3}{2}c^2(m-2)+9c^2\sigma_m\sqrt{m'}, \left(3k-\frac{3}{2}\right)a^2-\frac{1}{2}c^2(m-2)+3c^2\sigma_m\sqrt{m'}$
85 $\frac{2ia\sigma\sigma_m\left(\sqrt{k}\sigma_k\mathrm{sn}(\xi\|k)^2+1\right)\left(\sqrt[4]{m'}\sigma_m^{3/2}+\mathrm{dn}(\eta\|m)\right)}{c\left(\sigma_m\sqrt{m'}-1\right)\mathrm{sn}(\xi\|k)\left(\mathrm{dn}(\eta\|m)-\sigma_m^{3/2}\sqrt[4]{m'}\right)}$	$\frac{64\sqrt{k}\sigma_k\left(\sigma_k+\sqrt{k}\right)^2}{\left(\sigma_m\sqrt{m'}-1\right)^4}, (k+1)a^2+6\sqrt{k}\sigma_k a^2-\frac{3}{2}c^2(m-2)+9c^2\sigma_m\sqrt{m'}, 3(k+1)a^2+18\sqrt{k}\sigma_k a^2-\frac{1}{2}c^2(m-2)+3c^2\sigma_m\sqrt{m'}$
86 $\frac{2a\,\sigma\sigma_k\sigma_m\sqrt{k}\left(k+\sigma_k\sqrt{-kk'}\right)\left(\mathrm{cn}(\xi\|k)-\frac{\sigma_k\sqrt{-\frac{k'}{k}}}{\mathrm{cn}(\xi\|k)}\right)}{c\left(\sigma_m\sqrt{m'}-1\right)}$ $\times\frac{\left(\sigma_m^{3/2}\sqrt[4]{m'}+\mathrm{dn}(\eta\|m)\right)}{\left(\sigma_m^{3/2}\sqrt[4]{m'}-\mathrm{dn}(\eta\|m)\right)}$	$-\frac{64\sigma_k\left(\sqrt{-k'}-\sqrt{k}\sigma_k\right)^2\sqrt{-kk'}}{\left(\sigma_m\sqrt{m'}-1\right)^4}, (1-2k)a^2+6\sigma_k\sqrt{-kk'}a^2-\frac{3}{2}c^2(m-2)+9c^2\sigma_m\sqrt{m'}, (3-6k)a^2+18\sigma_k\sqrt{-kk'}a^2-\frac{1}{2}c^2(m-2)+3c^2\sigma_m\sqrt{m'}$
87 $\frac{2a\sigma\sigma_k\sigma_m\left(\sqrt{k'}-\mathrm{dn}(\xi\|k)^2\sigma_k\right)\left(\sqrt[4]{m'}\sigma_m^{3/2}+\mathrm{dn}(\eta\|m)\right)}{c\left(\sigma_m\sqrt{m'}-1\right)\mathrm{dn}(\xi\|k)\left(\mathrm{dn}(\eta\|m)-\sigma_m^{3/2}\sqrt[4]{m'}\right)}$	$-\frac{64\sigma_k\left(\sqrt{k'}-\sigma_k\right)^2\sqrt{k'}}{\left(\sigma_m\sqrt{m'}-1\right)^4}, (k-2)a^2+6\sigma_k\sqrt{k'}a^2-\frac{3}{2}c^2(m-2)+9c^2\sigma_m\sqrt{m'}, 3(k-2)a^2+18\sigma_k\sqrt{k'}a^2-\frac{1}{2}c^2(m-2)+3c^2\sigma_m\sqrt{m'}$
88 $\frac{a\sigma\left(\sigma_m\sqrt{k}-\sigma_k\right)\left(\sigma_k^{3/2}+\sqrt[4]{k}\mathrm{sn}(\xi\|k)\right)}{c\left(\sigma_m\sqrt{m'}-1\right)\left(\sigma_k^{3/2}-\sqrt[4]{k}\mathrm{sn}(\xi\|k)\right)}$ $\times\frac{\left(\sigma_m^{3/2}\sqrt[4]{m'}+\mathrm{dn}(\eta\|m)\right)}{\left(\sigma_m^{3/2}\sqrt[4]{m'}-\mathrm{dn}(\eta\|m)\right)}$	$\frac{\left(\sqrt{k}\sigma_k-1\right)^4}{\left(\sigma_m\sqrt{m'}-1\right)^4}, \frac{1}{2}(-k-1)a^2-3\sqrt{k}\sigma_k a^2-\frac{3}{2}c^2(m-2)+9c^2\sigma_m\sqrt{m'}, -\frac{3}{2}(k+1)a^2-9\sqrt{k}\sigma_k a^2-\frac{1}{2}c^2(m-2)+3c^2\sigma_m\sqrt{m'}$

continue I

number 2 $\sqrt{\nu}\frac{\partial\tan^{-1}}{\partial x}$		$\frac{c^4}{a^4}, \frac{b}{a}, \frac{d}{c}$
89	$\frac{a\sigma\sigma_m\left(\sqrt{k'}\sigma_k+\sqrt{-k}\right)\left(\sqrt[4]{k'}\left(-\sigma_k\right)^{3/2}+\sqrt[4]{-k}\mathrm{cn}(\xi\|k)\right)}{c\left(\sigma_m\sqrt{m'}-1\right)\left(\sqrt[4]{-k}\mathrm{cn}(\xi\|k)-\left(-\sigma_k\right)^{3/2}\sqrt[4]{k'}\right)}$ $\times\frac{\left(\sigma_m^{3/2}\sqrt[4]{m'}+\mathrm{dn}(\eta\|m)\right)}{\left(\sigma_m^{3/2}\sqrt[4]{m'}-\mathrm{dn}(\eta\|m)\right)}$	$\frac{\left(\sqrt{k'}\sigma_k+\sqrt{-k}\right)^4}{\left(\sigma_m\sqrt{m'}-1\right)^4}, \left(k-\frac{1}{2}\right)a^2 + 3\sigma_k\sqrt{-kk'}a^2 - \frac{3}{2}c^2(m-2) + 9c^2\sigma_m\sqrt{m'}, \left(3k-\frac{3}{2}\right)a^2 + 9\sigma_k\sqrt{-kk'}a^2 - \frac{1}{2}c^2(m-2) + 3c^2\sigma_m\sqrt{m'}$
90	$\frac{ia\sigma\sigma_m\left(\sqrt{k'}\sigma_k+\sqrt{-k}\right)\left(\sqrt[4]{k'}\sigma_k^{3/2}+\mathrm{dn}(\xi\|k)\right)}{c\left(\sigma_m\sqrt{m'}-1\right)\left(\mathrm{dn}(\xi\|k)-\sigma_k^{3/2}\sqrt[4]{k'}\right)}$ $\times\frac{\left(\sigma_m^{3/2}\sqrt[4]{m'}+\mathrm{dn}(\eta\|m)\right)}{\left(\sigma_m^{3/2}\sqrt[4]{m'}-\mathrm{dn}(\eta\|m)\right)}$	$\frac{\left(\sigma_k\sqrt{k'}-1\right)^4}{\left(\sigma_m\sqrt{m'}-1\right)^4}, -\frac{1}{2}(k-2)a^2 + 3\sigma_k\sqrt{k'}a^2 - \frac{3}{2}c^2(m-2) + 9c^2\sigma_m\sqrt{m'}, -\frac{3}{2}(k-2)a^2 + 9\sigma_k\sqrt{k'}a^2 - \frac{1}{2}c^2(m-2) + 3c^2\sigma_m\sqrt{m'}$

Note: $\theta[x]$ is θ function,it satisfies $x<0,\ \theta[x]=0; x\geq 0,\ \theta[x]=1.\ \sigma_m^2=1,\ \sigma_k^2=1.$

IV. CONCLUSION

So we can conclude that If two Jacobi elliptic function solutions of a NDE satisfy (1) and (2), the two solutions are equivalent. According to the usage of the modulus and phase transformation of elliptic functions to mKdV, a lot of the eqivalent Jacobi elliptic function solutions can be obtained when we knowing one of the Jacobi elliptic function solutions to some NDEs. It can be used to test the eqivalent of the other NDEs' Jacobi elliptic function solutions.

Appendix: Definition and Jacobi elliptic function

Common Jacobi elliptic function have three kinds[6,7] :sn($x|k$), cn($x|k$) and dn($x|k$), where k is the module of the Jacobi elliptic function. They meet the relations

$$\mathrm{sn}(x|k)^2+\mathrm{cn}(x|k)^2=1, \mathrm{dn}(x|k)^2+k\mathrm{sn}(x|k)^2=1,\ \mathrm{dn}(x|k)^2-k\mathrm{cn}(x|k)^2=1-k=k'. \tag{66}$$

It is known that

$$\mathrm{sn}(-x|k)=-\mathrm{sn}(x|k),\ \mathrm{cn}(-x|k)=\mathrm{cn}(x|k),\ \mathrm{dn}(-x|k)=\mathrm{dn}(x|k), \tag{67}$$

where k' is complementary module of Jacobi elliptic function. Except sn($x|k$), cn($x|k$) and dn($x|k$), other nine Jacobi elliptic functions can be defined

$1/\mathrm{sn}(x|k)\equiv\mathrm{ns}(x|k),\ \mathrm{dn}(x|k)/\mathrm{cn}(x|k)\equiv\mathrm{dc}(x|k),\ \mathrm{cn}(x|k)/\mathrm{dn}(x|k)\equiv\mathrm{cd}(x|k),$
$1/\mathrm{cn}(x|k)\equiv\mathrm{nc}(x|k),\ \mathrm{dn}(x|k)/\mathrm{sn}(x|k)\equiv\mathrm{ds}(x|k),\ \mathrm{sn}(x|k)/\mathrm{dn}(x|k)\equiv\mathrm{sd}(x|k),$
$1/\mathrm{dn}(x|k)\equiv\mathrm{nd}(x|k),\ \mathrm{sn}(x|k)/\mathrm{cn}(x|k)\equiv\mathrm{sc}(x|k),\ \mathrm{cn}(x|k)/\mathrm{sn}(x|k))\equiv\mathrm{cs}(x|k).$

Jacobi elliptic function has some common phase transformation, such as

$$\mathrm{sn}(x+2pK+2iqK'|k)=(-1)^p\mathrm{sn}(x|k), \tag{68}$$

$$\mathrm{cn}(x+2pK+2iqK'|k)=(-1)^{p+q}\mathrm{cn}(x|k), \tag{69}$$

$$\mathrm{dn}(x+2pK+2iqK'|k)=(-1)^q\mathrm{dn}(x|k), \tag{70}$$

$$\mathrm{sn}(x+K+iK'|k)=\frac{\mathrm{dc}(x|k)}{\sqrt{k}}, \tag{71}$$

$$\mathrm{cn}(x+K+iK'|k)=-i\sqrt{k'/\ k}\mathrm{nc}(x|k), \tag{72}$$

$$\mathrm{dn}(x+K+iK'|k)=-i\sqrt{k'}\mathrm{sc}(x|k), \tag{73}$$

$$\mathrm{sn}(x+K|k)=\mathrm{cd}(x|k), \tag{74}$$

$$\mathrm{cn}(x+K|k) = \sqrt{k'}\,\mathrm{sd}(x|k), \tag{75}$$

$$\mathrm{dn}(x+K|k) = \sqrt{k'}\,\mathrm{nd}(x|k), \tag{76}$$

$$\mathrm{sn}(x+iK'|k) = \frac{1}{\sqrt{k}}\mathrm{ns}(x|k), \tag{77}$$

$$\mathrm{cn}(x+iK'|k) = \frac{-i}{\sqrt{k}}\mathrm{ds}(x|k), \tag{78}$$

$$\mathrm{dn}(x+iK'|k) = \mathrm{cs}(x|k), \tag{79}$$

where $i^2 = -1$, $K = \int_0^{\pi/2} [1 - k\sin^2(\theta)]^{-\frac{1}{2}} d\theta$, $K' = \int_0^{\pi/2} [1 - k'\sin^2(\theta)]^{-\frac{1}{2}} d\theta$, p and q are integers.

[1] Liu S K, Fu J T, Liu S D, et al. Jacobi elliptic function expansion method and periodic wave solutions of nonlinear wave equations[J]. Phys. Lett. A, 2001, 289(1-2):69-74.

[2] Fu Z T, Liu S K, Liu S D, et al. New Jacobi elliptic function expansion and new periodic solutions of nonlinear wave equations[J]. Phys. Lett. A, 2001, 290(1-2):72-76.

[3] Liu G T, Fan T Y. New applications of developed Jacobi elliptic function expansion methods J][J]. phys. Lett. A, 2005, 345(1):161-166.

[4] Shen S F, Pan Z L, A note on the Jacobi elliptic function expansion method[J]. Phys. Lett. A, 2001, 308(1):143-148.

[5] Kevrekidis P G, Khare A, Saxena A and Herring G. On Some Classes of mKdV Periodic Solutions[J]. J. Phys. A, Math. Gen, 2004, 37(45):10959-10967.

[6] Gao B Q Elliptic Functions and Applications[M] (Beijing National Defence Industry Press, 1991).

[7] Mathematics Handbook Writing Group, Math Handbook(Higher Education Press, 2006).

[8] Dixon A C. The elementary properties of elliptic functions with examples[M], London:Macmillan, 1894.

[9] Kevrekidis P G, AKhare, A Saxena. Breather lattice and its stabilization for the modified Korteweg-deVries equation[J]. Phys. Rev. E, 2003, 68(4):047701-047704.

[10] Yan Z Y. New binary travelling-wave periodic solutions for the modified KdV equation[J]. Physics Letters A, 2008, 372(7):969-977.

[11] Fu Z T, Liu S D, Liu S K. A systematical way to find breather lattice solutions to the positive mKdV equation[J]. Journal of Physics. A, 2005, 40(18):4739-4746.